

DIRECTIONS FOR USE
INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated area and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites and cadellies: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surfaces application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

EPA REG. No. 6020-11
EPA EST. 6020-FI-1

FAST ACTION NO STAIN

ZAP

The Space Age Insecticide That Does the Job

ACTIVE INGREDIENTS:

Pyrethrins.	0.1%
*Piperonyl Butoxide, technical.	1.0%
Petroleum distillate.	0.4%
INERT INGREDIENTS.	98.5%

*Equivalent to 0.3% (butylcarbityl) (6-propylpiperonyl) ether and to 0.2% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Net Contents ____ Gallon

SPACE SPRAYING: Remove pets, birds and cover fish aquaria before spraying. Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths: direct spray at an upward angle, distributing it uniformly throughout the entire area, at a rate of 3-6 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited. Apply at a rate of 1½ gallons per acre (4½ ounces per 1000 square feet).

CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

DIRECTIONS FOR USE

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated area and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites and cadellies: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surfaces application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested carpeted beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Thoroughly liberally treat aptly, chests, closets, bureaus, and other storage areas directing the spray into cracks, joints, and crevices. After removal of infested articles may be treated lightly keeping nozzle at least 12 inches from fabric to avoid staining.

FOR FURTHER INFORMATION
CONTACT YOUR LOCAL
SPRAYER DEALER

FAST

ACTION

NO
STAIN

ZAP

The Space Age Insecticide That Does the Job

ACTIVE INGREDIENTS

Pyrethrins.....

*Piperonyl Butoxide.....

Petroleum.....

INERT INGREDIENTS

*Equivalent to 0.2%

KEEP OUT OF REACH OF CHILDREN

USE ONLY AS DIRECTED

SPACE SPRAYING: Remove all food and dishes from the area before spraying. Close doors and windows and shut off ventilation systems. Apply with a good marginator and fogger or fog machine or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths direct spray at an upward angle, distributing it uniformly throughout the entire area, at a rate of 3-6 ounces per 1000 cubic feet of space.

FOGGING: Fog in exposed or accessible stages of infestation of stored foodstuffs, at a rate of one gallon to 20,000 cubic feet of space. Contact insects directly whenever possible.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited. Apply at a rate of 1½ gallons per acre (4½ ounces per 1000 square feet).

CAUTION

Harmful if swallowed. Avoid contamination of feed and food.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of fish ponds. Do not apply where runoff is likely to occur. Do not apply in weather conditions favor drift from areas of use. Do not contaminate water by cleaning of equipment or disposal of excess spray. Use product only as specified on this label.

Do not reuse empty container.
Keep in a safe place.