




GUARANTEE

As per Interim DY-NAMO, contains Phenol Coefficient of 16 against *Staphylococcus aureus* ATCC 6538 and a Phenol Coefficient of 13.3 against *Salmonella typhosa* ATCC 6539 when tested by the A.O.A.C. Official Method, 11th Edition, 1955.

DISINFECTANT	BACTERICIDE
CLEANER	VIRUCIDE**
DEODORIZER	PATHOGENIC FUNGICIDE
	REDUCES SURFACE TENSION
	RESIDUAL BACTERIOSTAT

* Kills mildew fungi on contact
 ** DY-NAMO is an effective Virucide for Influenza Virus Type A2 (Hong Kong) on inanimate environmental surfaces.

GENERAL INFORMATION

DY-NAMO contains an effective synthetic detergent system and builders for removal of general industrial soils such as would normally be encountered in hospitals, nursing homes, schools and industrial plant restrooms. DY-NAMO also reduces the surface tension of water, improving its wettability so that the disinfectant can reach crevices in subvisible pores, scratches and cracks of surfaces being treated. After the initial demand for disinfectant has been satisfied, the remaining DY-NAMO will establish a residual bacteriostatic action.

By the following test methods, DY-NAMO will kill the following organisms on inanimate surfaces at recommended use dilutions.

PATHOGENIC BACTERIA:	FUNGI:
<i>Escherichia coli</i>	<i>Aspergillus niger</i>
<i>Staphylococcus aureus</i>	<i>Candida albicans</i>
<i>Salmonella typhosa</i>	<i>Trichophyton interdigitale</i>
<i>Shigella flexneri</i>	
<i>Shigella sonnei</i>	
<i>Shigella dysenteriae</i>	
<i>Shigella flexneri</i>	

Net Contents

GALLONS

DY-NAMO

DISINFECTANT - DETERGENT - DEODORANT

Phenol Coefficient (16) - *Staphylococcus aureus* ATCC 6538

Phenol Coefficient (13.3) - *Salmonella typhosa* ATCC 6539

ACTIVE INGREDIENTS

Alkyl (C14, 90%, C12, 5%, C16, 5%) dimethyl dichloro-benzyl ammonium chloride	1.250%
Alkyl (C14, 5%, C16, 28%, C12, 14%) dimethyl benzyl ammonium chloride	0.625%
Alkyl (C14, 90%, C12, 5%, C16, 5%) dimethyl ethyl ammonium bromide	0.625%
Sodium carbonate	0.500%
Sodium metasilicate	0.500%
Essential oils	0.250%
Ethylenediamine, tetrasodium salt	0.190%

TOTAL ACTIVE	3.940%
INERT INGREDIENTS	96.060%

TOTAL INGREDIENTS 100.000%

** Contains sodium nitrate

WARNING

KEEP OUT OF REACH OF CHILDREN

(See side panel for First Aid and other precautions)

EPA Reg. No. 6020-11

MANUFACTURED BY

USE DIRECTIONS

FOR RESTROOMS AND GENERAL DISINFECTION IN HOSPITALS, NURSING HOMES, INDUSTRIAL AND INSTITUTIONAL USES. Wherever practical, rinse away surface soils with water. For cleaning, use 4 ounces DY-NAMO per gallon of water and thoroughly wet the surface to be cleaned. Cleaning may be accomplished with a clean mop, sponge, squeegee, wet vacuum system, or pressure sprayer. For disinfecting surfaces difficult to clean or contagious disease areas, use 8 ounces per gallon of water, thoroughly wetting all surfaces. Any food contact surfaces treated with DY-NAMO should be thoroughly rinsed with potable water before re-use.

FOR DISINFECTING AND DEODORIZING OF GARBAGE CANS, GARBAGE TRUCKS, INDUSTRIAL WASTE RECEPTACLES AND OTHER GARBAGE HANDLING EQUIPMENT. It is especially important to pre-clean, preferably using a pressure sprayer, in order for the disinfectant to perform properly. Then, apply a wetting solution of 8 ounces of DY-NAMO per gallon of water.

FOR HAND CLEANING OR PRESSURE CLEANING OF WALLS, FLOORS AND INDUSTRIAL WORK AREAS. Use 4 ounces per gallon of water. The use of rubber gloves is recommended for personnel doing continuous hand cleaning.

DY-NAMO may be used through automatic floor scrubbing equipment.

WARNING

KEEP OUT OF REACH OF CHILDREN. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. Avoid breathing spray mist.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of water or milk, or white. If these are not available, drink water. Do not induce vomiting. Avoid alcohol. Call a physician immediately.

Rinse empty container thoroughly with water before re-use.

EPA Est. No. 6020-FI-01