STOKER MALATHION DUST No. 5

Active Ingredient	
Malathion (O O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	5%
Inert Ingredients	95%
	100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

AVOID BREATHING DUST
AVOID REPEATED CONTACT WITH SKIN.
WASH THOROUGHLY AFTER USING
AVOID CONTAMINATION OF FEED AND FOOD

NOTE TO PHYSICIANS: Malathion when repeated, prolonged and extensive use may cause cholinesterase inhibition. Atropine is antidotal.

DIRECTIONS FOR USE

Apply [illegible] amounts of this product uniformly to the [illegible] when [illegible] is [illegible] Apply at the first sign of infestation and repeat at [illegible] intervals as needed. Maximum control can be observed when [illegible] for specific crops.

USE PRECAUTIONS

To [illegible] Do not contaminate water, food or feed by storage or disposal.

[illegible]

RECOMMENDATIONS

Alfalfa: Aphids, weevil, grass [illegible] Apply 25-30 lbs per acre. [illegible] Apply [illegible] Do not graze or feed treated forage within 7 days of last application. Do not apply within 24 hours of bloom.

Note: May be used [illegible]

BEANS: Aphids, cucumber beetle, Japanese beetle, Mexican bean beetle, [illegible] Apply 25-30 lbs per acre. Make [illegible] Do not apply within 24 hours of bloom.

CARROTS, RADISHES, SALSIFY, HORSERADISH: Aphids, [illegible] Apply [illegible] Do not apply within 7 days of harvest.

CLOVER: Aphids, spotted mites. Apply 25-30 lbs per acre. [illegible] Do not apply within 7 days of harvest. Do not feed or graze treated forage within 24 hours of application. Should be applied on day of harvest or pasturing.

COLE CROPS: Broccoli, Brussels sprouts, cabbage, cauliflower, kale, collards, kohlrabi. Aphids, [illegible] flea beetle, imported cabbage worm, cabbage looper, cabbage worm. Apply 25-30 lbs per acre. Make [illegible] application, repeating as necessary. Do not apply within 7 days of harvest. Brussels sprouts, cabbage, cauliflower, kale and mustard: 3 days. Broccoli or kohlrabi: 14 days.

COTTON: Aphids, boll weevil, [illegible] cotton bollworm, pink bollworm, saltmarsh caterpillar, cotton leafworm, [illegible] Fleahoppers, [illegible] bugs, stink bugs, tarnished plant bug, thrips, Lygus bugs, white flies. Apply 10-20 lbs per acre. Boll weevil, [illegible] armyworms, granulate cutworm, and garden webworms. Apply 20-40 lbs per acre. Make repeat applications as needed.

CUCUMBERS, SQUASH AND MELONS: Aphids, [illegible] and spider mites. Apply [illegible] lbs per acre. For cucumber beetles and squash vine borer use 35 lbs per acre. Apply when [illegible] and dust at base of plants. [illegible] CAUTION: Do not [illegible] Make no applications within [illegible] days of harvest.

GRAPES: Cutworms, grasshoppers, thrips, [illegible] Apply when [illegible] Make [illegible] applications as needed. Do not apply within 7 days of harvest.

LETTUCE: Aphids, cabbage looper, cutworms, mites. Apply 25-30 lbs per acre. [illegible] Do not allow animals to feed on treated crop within 24 hours of application. Do not apply within 7 days of harvest.

PEAS: Pea aphid, [illegible] weevil. Apply 25-30 lbs per acre. Make [illegible] Do not apply within 7 days of harvest. Do not feed or graze treated forage within 24 hours of application.

POTATOES: False chinch bug, [illegible] Apply 25-30 lbs per acre. For [illegible] potatoes, [illegible] apply 25-30 lbs per acre.

SPINACH: Aphids. Apply 25-30 lbs per acre. [illegible] Do not apply within 7 days of harvest.

STRAWBERRIES: Aphids, [illegible] Apply when [illegible] appear. For [illegible] leaf roller, strawberry root weevil, [illegible] white grubs. Apply 20-40 lbs per acre. For strawberry [illegible] weevil, [illegible] Make [illegible] Do not apply within 7 days of harvest. Do not feed or graze treated forage within 24 hours of application.

TOMATOES: Aphids, [illegible] Apply 25-30 lbs per acre. Make [illegible] Do not apply within 7 days of harvest.

ONIONS: Onion [illegible] maggots. Apply 25-30 lbs per acre. [illegible] Use first application when [illegible] Repeat every 7 days as [illegible] Do not apply within 7 days of harvest.

PASTURE & RANGE GRASS GRAIN & HAY: Aphids, [illegible] Apply 25-30 lbs per acre. Make [illegible] Do not graze or feed treated forage within 24 hours of application. Do not apply within 7 days of harvest.

Manufactured by

PRACTICAL PEST CONTROL
STOKER CO.

STOKER CO.

El Centro, California — Brawley, California

ACCEPTED

[illegible]