

STOKER DUSTING SULFUR

Active Ingredient:

Sulfur 98%

Inert

Ingredients 2%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Sulfur will cause severe fruit and leaf injury to sulfur sensitive crops. Do not apply to apricots, comice pears, cranberries, cucurbits (cucumber, cantaloupes, melons, squash), walnuts, or to other sensitive plants.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90° F.

Do not use sulfur with oil sprays or within 30 days of an oil spray.

Do not use sulfur on any crop unless local use has proved safe.

If the crop to be treated will be used for coming or freezing purposes, consult food processor before applying sulfur.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5 to 10 day intervals, depending on severity of infection, but observe use limits given for specific crops. Apply at the rate of 30 to 40 lb. unless otherwise specified.

RECOMMENDATIONS

ALFALFA: Spider mites - Apply 15-50 lbs. per acre. Caution: Sulfur may cause petal burn to open flowers.

BEANS: Powdery mildew, rust, spider mites - Apply 15-30 lbs. per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until bloom time. Caution: May cause injury to some varieties of beans.

CITRUS: Six-spotted and citrus rust mites - Apply 80 lbs. per acre (10 to 150 lb. per tree, depending on size, for citrus rust mite in Florida). Citrus flat and Yuma spider mites - Apply 100 lbs. per acre; citrus thrips - Apply 100 lbs. per acre in three applications: mid-March, petal fall, and 2-4 weeks later. Do not apply within 3 to 6 weeks of an oil spray.

GRAPES (VINEFERA VARIETIES): Rust, powdery mildew - Apply 15-20 lbs. per acre in 4 applications: when shoots are 4" long, 12-14" long, 14 days afterwards, 12-14 days thereafter to harvest. Sulfur is extremely phytotoxic to foliage of Concord and most other labrusca type grapes and should not be used on these varieties.

COTTON: Atlantic mite, cotton rust - Apply 25-30 lbs. per acre.

SAFFLOWER-SORGHUM-SUGAR BEETS: Spider mite - Apply 15 to 50 lbs. per acre. No time limitation.

MELON: Honeydew and sulfur-resistant melon only. Powdery mildew, spider mites - Apply 20-50 lbs. per acre.

PEAS: Powdery mildew - Apply 15-20 lbs. per acre.

STRAWBERRIES: Powdery mildew - Apply 15-20 lbs. per acre.

TOMATOES: Tomato russet mite - Apply 15-40 lbs. per acre.

ORNAMENTALS: Make full coverage applications on both sides of leaves. Keep new growth protected. Chrysanthemum: Powdery mildew, rust. Do not apply when flowering. Hollyhock: Rust, Rose: Black spot, powdery mildew, rust. Caution: Some burning may occur when used during warm weather.

NOTICE: Our recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of STOKER CO., no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe and sound practices. THE BUYER must assume all responsibility, including injury or damage resulting from its use as such or in combination with other products.

Net Weight 50 Pounds



For Best Results

STOKER CO.

El Centro, California - Brawley, California

