

STOKER PHOSDRIN* 400

NET CONTENTS

GALLONS

INGREDIENTS

ACTIVE INGREDIENTS:

- Phosphorus pentasulfide
- Phosphorus pentoxide
- Inert ingredients

*Equivalent to 4 lbs. PHOSDRIN
(Contains 4 lbs. active ingredient per gallon)

WARNING

FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.

DANGER

KEEP OUT OF REACH OF CHILDREN



POISON



SORGHUM (for forage and grain): Aphids, corn earworm, webworms, fall armyworms, etc. - 1 pt. For fall armyworms use 1 pt. dosage. Do not apply within 3 days of harvest. Do not apply Phosdrin to or allow spray to drift onto varieties of Sorghum which are sensitive to phosphates. To insure no result in burning of the Sorghum for Mill. Consult State Agricultural Extension Service prior to application.


SUMMER SQUASH: For control of aphids use 1/2 pt. per acre. For control of cabbage looper, dipterous leafminer (adults), leathoppers, big blight, nates, false chinch bugs, saltmarsh caterpillar, cutworm (larvina) and grasshoppers use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest.

MELONS (INCLUDING CANTALOUPE, HONEY-DEW MELONS, MUSKMELONS, WATERMELONS): For control of aphids use 1/2 pt. per acre. For control of Cabbage looper, dipterous leafminer (adult), leathoppers, lycus bugs, nates, false chinch bugs, saltmarsh caterpillars, cutworms (larvina) and grasshoppers use 1/2 to 1 pt. per acre. To control rim worms, cabbage looper, cutworms, saltmarsh caterpillars, tobacco hornworms on watermelons apply 1/2 pt. per acre. Do not treat cantaloupes, honey-dew melons, or watermelons within 1 day of harvest.

STOKER CO.
BRAWLEY, CALIFORNIA
EPA REG. NO. 06023-00033AA

STOKER CO.
BRAWLEY, CALIFORNIA
EPA REG. NO. 06023-00033AA

STOKER CO.



Brawley, California

EPA Reg. No. 06023-00033AA

ACCEPTED WITH COMMENTS

USE AND DIRECTIONS FOR APPLICATION

ALFALFA, CLOVER: Aphids, alfalfa caterpillar, alfalfa caterpillar, alfalfa seed fly, alfalfa caterpillar, alfalfa caterpillar, grasshoppers, leafhoppers, lygus bugs, mites, 1 - 1 pt. Do not apply within 1 day of harvest.

BEANS: Aphids, grasshoppers, leafhoppers, Mexican bean beetle, mites, 1 - 1 pt. Do not apply within 1 day of harvest.

BRUSSELS SPROUTS, CAULIFLOWER: Aphids, cabbage looper, climbing cutworm, grasshopper, leafhopper, lygus bug, mite, saltmarsh caterpillar, apply 1-2 pt. Do not apply within 3 days of harvest.

BROCCOLI, CABBAGE: For control of aphids use 1/2 to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, saltmarsh caterpillar, mites cutworms (climbing), dipterous leafminer (adult) and lygus bugs use 1/2 to 1 pt. per acre. Do not treat within 1 day of harvest. For hard to kill insects, including aphids, use 1 qt. per acre, but do not treat within 3 days of harvest.

CARROTS: Aphids, cabbage looper, climbing cutworms, dipterous leaf miner (adults), leafhoppers, lygus bugs, mites, salt-marsh caterpillar, 1 - 1 pt. Do not apply within 1 day of harvest.

CUCUMBER: Aphids, grasshoppers, leafhoppers, mites, 1 - 1 pt. Do not apply within 1 day of harvest.

LETTUCE: Aphids, Cabbage looper, corn earworm, climbing cutworm, adult dipterous leafminer, False chinchbug, grasshopper, imported cabbageworm, lygus bug, mite, salt-marsh caterpillar, thrips, 1 - 2 pt. or 1 - 1 pt, do not apply within 4 days of harvest at the 1 - 2 pt. per acre rate. Do not apply within 2 days of harvest at the 1 - 1 pt. per acre rate.

OKRA: Aphids, climbing cutworms, corn earworms, green stinkbug, mites, velvet bean caterpillar, 1 - 1 pt. Do not apply within 1 day of harvest.

ONIONS (including green onions): Climbing cutworms, thrips, 1 - 1 pt. Do not apply within 1 day of harvest.

POTATOES: Aphids, grasshoppers, leafhoppers, mites, 1/2 pt. per acre. Do not apply within 1 day of harvest.

SPINACH: Aphids, Cabbage looper, Climbing cutworm, adult dipterous leafminer, false chinchbug, grasshopper, leafhopper, imported cabbageworm, mite, saltmarsh caterpillar, 1 - 1 pt. Do not apply within 4 days of harvest.

TOMATOES: Aphids, grasshoppers, leafhoppers, mites, 1/2 - 1 pt. Do not apply within 1 day of harvest.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

FIRST AID TREATMENT: Atropine is the emergency antidote for Phosdrin poisoning. Atropine is obtainable only on a Doctor's Prescription. Consult your physician about obtaining an adequate supply of 1-100 grain Atropine Tablets for emergency use.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water) or soapy water. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. If material has been spilled on the skin, immediately remove the patient from the vicinity of the insecticide. Remove contaminated clothing and wash skin with soap and water. If the material has been splashed into the eyes, wash immediately with clear water for at least ten minutes. If warning symptoms appear, administer two 1-1000 grain Atropine Tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (SEE WARNING SYMPTOMS ABOVE). Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

2-PAM is also antidotal and may be administered in conjunction with atropine.

TREATMENT: Have victim lie down and keep quiet. Atropine is the specific therapeutic antagonist of choice against the parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, Atropine Sulfate should be injected at 10 minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. All warn further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

DISPOSAL: Rinse spray equipment and container. Bury liquid waste in non-crop lands located away from water supplies. Port rate empty metal containers, crush and bury glass.

NOTICE: The recommendations for the use of this product are based upon information believed to be reliable at time of printing. The use of this product being beyond the control of STOKER CO., no guarantee, expressed or implied is made as to the effects or results to be obtained if used in accordance with directions as established safe and sound practices. THE BUYER must assume all responsibility for injury or damage resulting from its misuse as such or in combination with other products.

ACCEPTED

1-22-74