

OIL-1-CIDE "99"

CITRUS SPRAY OIL

ANALYSIS

ACTIVE INGREDIENTS: Petroleum Oil (Min. U. R. 92%)	99.0%
INERT INGREDIENTS	1.0%

USDA REG. No. 6031-2

MANUFACTURED BY

INTERNATIONAL LUBRICANT CORP.

NEW ORLEANS, LOUISIANA, U. S. A.

DISTRIBUTED BY

NIAGARA CHEMICAL DIV.

F. M. C. CORPORATION

LOS FRESNOS, TEXAS


CAUTION: KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED.
SEE OTHER PRECAUTIONS ON THE BACK SIDE PANEL.


ACCEPTED

SEP 14 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTRATION
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

66517





**FISH AND/OR WILDLIFE CAUTION
DO NOT CONTAMINATE WATER BY CLEANING OF
EQUIPMENT, OR DISPOSAL OF WASTES**

NOTICE

The company guarantees this material to consist of the ingredients specified but makes no other representation or warranty, express or implied. Use of this material in a manner or at a time other than in accordance with the directions may produce plant injury. No representative of the company may vary any of the foregoing and the buyer hereby accepts said material subject to all of the terms hereof.

DIRECTIONS FOR USE ON CITRUS

OIL-I-CIDE-99 is a specially prepared Emulsifiable Citrus Spray Oil. If properly applied it will control the following insects which attack Citrus: California Red Scale, Florida Red Scale, Chaff Scale, Purple Scale, Long Scale, Brown Soft Scale, Cottony Cushion Scale, Red Spider Mites, and Whiteflies.

OIL-I-CIDE-99 should be mixed at the rate of 1½ gallons per 100 gallons of water which will give a 1.5% actual oil in the final solution. Thorough coverage of all parts of the tree is essential for proper control. High pressure high volume spray equipment should be used which is capable of pumping 20 gallons per minute at a pressure of at least 500 - 600 lbs.

CAUTION

OIL-I-CIDE-99 is compatible with most of the other insecticides, fungicides, and nutritional sprays commonly used on Citrus, such as Malathion, Zineb, and Copper compounds. DO NOT USE WITH SULPHUR OR WITHIN 30 DAYS BEFORE OR AFTER SULPHUR HAS BEEN USED. Do not use OIL-I-CIDE-99 when temperature is above 90°F or below 40°F. Do not use OIL-I-CIDE-99 on bearing trees after September 1st since application after this date may interfere with coloring of early fruit and may make the trees more susceptible to cold injury. OIL-I-CIDE-99 should not be applied to drought weakened trees.

CAUTION:

**KEEP OUT OF REACH OF CHILDREN.
HARMFUL IF SWALLOWED.**