

INSTRUCTIONS FOR USE

FOR RATS: Place 1 oz. to 1 lb. of WARFET in locations frequented by rats. Four ounces of bait per bait location should be the minimum amount placed, maintaining an uninterrupted supply for approximately 10 days. Protect from children, dogs, cats and livestock by means of bait boxes or cages where necessary. Baits should be replaced as consumed. Where a continuous source of infestation prevails from a dump or field, permanent bait stations should be used and baits replenished as needed.

For controlling mice, follow the same general directions as for rats except bait placements may be in tablespoon amounts and placed at 8-12 ft. intervals. An uninterrupted supply of bait should be maintained for at least 15 days. Mice are more difficult to control than rats.

*The addition of T.M.F. to Warfarin is an exclusive development by Vineland Laboratories, Inc.

Net weight of

Warfarin (50% active) 15.000


Warfarin (50% active) 15.000

Warfarin (50% active)

Warfarin (50% active) 15.000

Warfarin (50% active) 15.000

Warfarin (50% active) 15.000



VINELAB®

WARFET

(Trademark)

Ready-To-Use Rat and Mouse



**New, Improved Warfarin R
With T.M.F. Added
For Hard-to-Kill Rats and**

CAUTION: Keep Out of Reach

WARNING: See Right Panel For Ad

Net Weight (Av.) **5 POUNDS**

**VINELAND LABORATORIES
VINELAND, NEW JERSEY**

VINELAB®

WARFET

Trademark

To-Use Rat and Mouse Bait



**Improved Warfarin Rodenticide
With T.M.F. Added
Hard-to-Kill Rats and Mice**

Keep Out of Reach of Children

See Right Panel For Additional Cautions

Net Weight (Av.) 5 POUNDS

**AND LABORATORIES, INC.
VINELAND, NEW JERSEY**

IMPORTANT

Bait should be placed in locations not accessible to children, pets, and domestic animals or in tamper-proof bait boxes.

CAUTION: WARFET contains as its active ingredient an anticoagulant chemical which, if taken by humans, domestic animals or pets, may increase the clotting time of the blood and hemorrhage may result. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for Warfarin. In case baits are accidentally eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS: When a human has been known to have ingested WARFET, blood transfusions combined with intravenous injections and oral doses of vitamin K are indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin.

U.S.D.A. Reg. No. 6035-25

MADE IN U.S.A.

REF. 69-00