

DIRECTIONS

Thoroughly mix 1 part RATRID with 19 parts of fresh bait material such as corn meal. Bait made with fresh ground meat, fish or vegetables should be set in late afternoon and replaced daily with fresh bait. Where a continuous source of infestation is present (nearby dumps) permanent bait stations should be established and replenished as needed.

For Rats: Four ounces of bait per bait location where rats are known to frequent should be the minimum amount used. An uninterrupted supply of bait should be maintained for at least 10 days. Death results in 5 to 14 days.

For Mice: Mice are more difficult to control than rats, making it necessary to have more bait stations. Tablespoon amounts (1/4 to 1/2 ounce) of bait per location should be placed at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

IMPORTANT: Bait should be placed in areas inaccessible to children, pets, and other animals, and in areas of protection.

VINELAB®

RATRID

Rodenticide Concentrate

NET WEIGHT



1 POUND (AV.)

Made With

WARFARIN* plus T. M. F.®

For Hard-to-Kill Rats and Mice

CAUTION: Keep Out of Reach of Children

CAUTION: See side panel for additional precautions.

VINELAND LABORATORIES, INC.

Vineland, New Jersey

2-27-67
6035-26

VINELAB®

RATRID

Rodenticide Concentrate

NET WEIGHT



1 POUND (AV.)

Made With

WARFARIN* plus T. M. F.®

For Hard-to-Kill Rats and Mice

CAUTION: Keep Out of Reach of Children

CAUTION: See side panel for additional precautions.

VINELAND LABORATORIES, INC.
Vineland, New Jersey

CAUTION

If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. Dogs and cats will actively search out baits of this type. Therefore, baits must be placed so as to prevent feeding by these animals. It also contains an antibacterial agent which inhibits bacteria that produce Vitamin K, an antidote for Warfarin. In such case, intravenous and oral administration of Vitamin K with blood transfusions are indicated as in the case of hemorrhage caused by overdose of dihydroxycoumarin.

I.M.F. is an exclusive Vinelab process that tends to increase the palatability of food materials that are mixed with this concentrate.

Active ingredients:

•Warfarin (3-acetyl-4-hydroxycoumarin)	0.5
N-(2-quinoxalyl)-sulfamamide	0.5
•Sulfquinoxaline	0.5
Inert ingredients	99.0
Total	100.0

U.S.D.A. Reg. No. 6035-26

Made in U.S.A. R2-76-140

27-67
035-26