

EPA Est. No. F-696 NJ-2
EPA Reg. No. 627-A-37

Precautionary Statements

Hazards to Humans and Domestic Animals

The following statements apply to both humans and domestic animals:

If formaldehyde gas is inhaled: Remove person from source of fume. At first signs of irritation, give 100% oxygen. If breathing is difficult, give respiration by mouth-to-mouth resuscitation. Call physician or hospital immediately.

If formaldehyde is taken internally: Practical Treatment: Give 1/2 cup of Plain Milk or 1/2 cup of Diluted Lemonade. If water is available, dilute 1/2 cup of water with 1/2 cup of plain Distilled Water. Take one-half cup of this mixture, swallow it, and drink another half cup of water. Do not induce vomiting.

In case of contact: Flush with plenty of water for several minutes.

Physical and Chemical Hazards

FIRE HAZARD: Use extreme care in handling. In the presence of an open flame, formaldehyde should be kept at all times to prevent this from happening.

Air: Cover the storage should be flushed with water.



FORMALDEGEN-F

U.S. Patent No. 2,992,612

With the application of heat, FORMALDEGEN F will liberate Formaldehyde gas plus a compound which will fluoresce under ultra violet light.

FORMALDEGEN F is to be used on hatching eggs either on the farm or at the hatchery. The inclusion of the compound which will fluoresce under ultra violet light is for the purpose of identifying at a later time the fact that such eggs have been fumigated with Formaldehyde gas.

AMOUNTS
100 POUNDS
100 EGGS
1000 EGGS

DANGER: Keep Out of Reach of Children



See First Aid and Additional Precautions on Left Panel

NET WEIGHT 22 POUNDS (1 KILOGRAM)



VINELAND LABORATORIES, INC.
A Division of Vincor
VINELAND, NEW JERSEY

4

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. Place the formaldehyde fumigant in a suitable

closed vessel containing the eggs to be fumigated.

2. Heat the vessel containing the formaldehyde fumigant extra moisture and heat when necessary.

3. After the fumigation, allow the fumigant to be fumigated.

4. Use 20 grams of Formaldegen F to treat 100 eggs.

5. Use 200 grams of Formaldegen F to treat 1000 eggs.

6. Use 2 kilograms of Formaldegen F to treat 10,000 eggs.

7. Use 20 kilograms of Formaldegen F to treat 100,000 eggs.

8. Use 200 kilograms of Formaldegen F to treat 1,000,000 eggs.

9. Use 2 kilograms of Formaldegen F to treat 10,000,000 eggs.

10. Use 20 kilograms of Formaldegen F to treat 100,000,000 eggs.

11. Use 200 kilograms of Formaldegen F to treat 1,000,000,000 eggs.

12. Use 2 kilograms of Formaldegen F to treat 10,000,000,000 eggs.

13. Use 20 kilograms of Formaldegen F to treat 100,000,000,000 eggs.

14. Use 200 kilograms of Formaldegen F to treat 1,000,000,000,000 eggs.

15. Use 2 kilograms of Formaldegen F to treat 10,000,000,000,000 eggs.

16. Use 20 kilograms of Formaldegen F to treat 100,000,000,000,000 eggs.

17. Use 200 kilograms of Formaldegen F to treat 1,000,000,000,000,000 eggs.

18. Use 2 kilograms of Formaldegen F to treat 10,000,000,000,000,000 eggs.

19. Use 20 kilograms of Formaldegen F to treat 100,000,000,000,000,000 eggs.

20. Use 200 kilograms of Formaldegen F to treat 1,000,000,000,000,000,000 eggs.

Storage & Disposal

Do not reuse container. Wash thoroughly with water and detergent. Discard in safe place. Do not use, pour, spill or store near heat or open flame.

30