

U.S.D.A. Reg. No. 6063-52

ONE POUND TREATS UP TO 4000 SQUARE FEET

THIS IS THE NEW ALL-SOLUBLE COMBINATION, CONTAINING ADDITIONAL STICKER, SPREADER AND ACTIVATOR.

DIRECTIONS

1. Be sure soil is moist and lawn healthy before treating. Do NOT apply to newly seeded lawns or to lawns stunted and dormant because of drought. During drought, water grass well several days in advance, and restore turf to healthy green color with Comar Lawn Food before treatment.

2. Keep lawn mowed to about 1 1/2" to 2" during treatment.

HOW TO APPLY: Use hose gun, or watering can: coarse sprays are less likely to drift.

For Common Broad-Leaf Weeds, and Young Seeding Crabgrass use 1 tablespoon per gallon of water applied to 65 square feet of lawn at temperatures above 85 F., and 1 1/2 tablespoons per gallon of water per 65 square feet for temperatures below 85 F.

On older or branched crabgrass, or hard-to-kill weeds, use 2 tablespoons per gallon of water in hot weather, per 65 square feet of lawn. If weather is cool (70-80 F), double dosage in each case. The interval between treatments should be one week.

Lawn may slightly discolor from treatment, but if directions are followed, this should be only temporary, and your good turf grasses will quickly return to normal color after first cutting.

In cool weather, weeds may take some time to show effect, but do not be impatient, since kill will result. Do not apply until grass is dry. Apply on a still day, to avoid drift. Do not water until a day after application. Repeat treatment as needed, at 2 to 3 week intervals. Flush spray jar and head thoroughly after use with mild ammonia solution (2 tablespoons per gallon).

WILL NOT INJURE TURF WHEN USED AS DIRECTED


TO SPRAY LARGE AREAS: Measure out 1000 square feet (20' x 50') and place required number of tablespoons in hose sprayer jar. Fill jar to top with water, dissolving powder. Apply evenly to area, until jar is empty.

- KILLS BROADLEAF WEEDS FROM ROOT TO CROWN
- KILLS CRABGRASS



CONCENTRATED • DOUBLE THREAT

CRABGRASS & WEED KILLER



ACTIVE INGREDIENTS:
 Disodium Methanearsonate, Anhydrous* 15.75%
 Sodium Salt of 2, 4-Dichlorophenoxyacetic Acid** 10.00%
 INERT INGREDIENTS 74.25%
 100.00%

*Total Arsenic as elemental, all in water soluble form, 6.42%, equivalent to 25.0% Disodium Methanearsonate, Hexahydrate.

**2, 4-Dichlorophenoxyacetic Acid equivalent 9.5% minimum.

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL

CAUTION

Keep tightly closed when not in use.

Avoid contact with eyes or clothing. May cause skin irritation. Avoid contact with skin or breathing of spray mist. Avoid contamination of feed or foodstuffs. Store beyond reach of children or domestic animals. Keep children and animals off treated areas until washed into soil. Wash hands after using. Harmful If Swallowed. Not to be taken internally.

ANTIDOTE: If taken internally, induce vomiting, and call a doctor at once.

Do not permit weed killer dust or spray to come too close to beneficial broad leaf plants, flowers, trees or crops. Do not store near seeds, insecticides, fungicides or fertilizer. Avoid spraying in wind. Vapors from this product may cause injury to nearby plants. Coarse sprays are less likely to drift.

Kills

MANY VARIETIES OF BROAD AND NARROW LEAVED WEEDS, SUCH AS:

CRABGRASS	BULL THISTLE	LAMB'S QUARTERS
DANDELION	BURDOCK	MORNING GLORY
BUCKHORN	CHARLOCK	WILD MUSTARD
PLANTAIN	CHICORY	PEPPER GRASS
BINDWEED	GIANT RAGWEED	PIGWEEN
TREFOIL	GOLDENROD	STINGING NETTLE

This spray is selective in the presence of Kentucky, Merion Blue, and Zoysia grasses.

DO NOT USE ON Dichondra, Centipede, Bermuda, St. Augustine grasses, or on lawns predominantly of Bent grass. This spray may injure clover.

COMAR INDUSTRIES
HANOVER, PA. 17331

DO NOT RE-USE CONTAINER, DESTROY WHEN EMPTY

Copr. Comar Industries, Incorporated
NET WEIGHT 1 LB.


U.S.D.A. Reg. No. 6063-52

Comar

CONCENTRATED • DOUBLE THREAT

CRABGRASS & WEED KILLER

ACCEPTED
4-7-71
6063-52



ACTIVE INGREDIENTS:
Disodium Methanesulfonate, Anhydrous* 15.75%
Sodium Salt of 2, 4-Dichlorophenoxyacetic Acid** 10.00%
INERT INGREDIENTS 74.25%
100.00%

*Total Arsenic as elemental, all in water soluble form, 6.42% equivalent to 25.0% Disodium Methanesulfonate, Hexahydrate.
**2, 4-Dichlorophenoxyacetic Acid equivalent 9.5% minimum.

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL

CAUTION

Keep tightly closed when not in use.

Avoid contact with eyes or clothing. May cause skin irritation. Avoid contact with skin or breathing of spray mist. Avoid contamination of feed or foodstuffs. Store beyond reach of children or domestic animals. Keep children and animals off treated areas until washed into soil. Wash hands after using. Harmful if Swallowed. Not to be taken internally.

ANTIDOTE: If taken internally, induce vomiting, and call a doctor at once.

Do not permit weed killer dust or spray to come too close to beneficial broad leaf plants, flowers, trees or crops. Do not store near seeds, insecticides, fungicides or fertilizer. Avoid spraying in wind. Vapors from this product may cause injury to nearby plants. Coarse sprays are less likely to drift.

Kills MANY VARIETIES OF BROAD AND NARROW LEAVED WEEDS, SUCH AS:

CRABGRASS	BULL THISTLE	LAMB'S QUARTERS
DANDELION	BURDOCK	MORNING GLORY
BUCKHORN	CHARLOCK	WILD MUSTARD
PLANTAIN	CHICORY	PEPPER GRASS
BINDWEED	GIANT RAGWEED	PIGWEE
TREFOIL	GOLDENROD	STINGING NETTLE

This spray is selective in the presence of Kentucky, Marion Blue, and Zoysia grasses.

DO NOT USE ON Dichondra, Centipede, Bermuda, St. Augustine grasses, or on lawns predominantly of Bent grass. This spray may injure clover.

COMAR INDUSTRIES
HANOVER, PA. 17331

DO NOT RE-USE CONTAINER, DESTROY WHEN EMPTY

Copr. Comar Industries, Incorporated
NET WEIGHT 1 LB.

ONE POUND TREATS UP TO 4000 SQUARE FEET

THIS IS THE NEW ALL-SOLUBLE COMBINATION, CONTAINING ADDITIONAL STICKER, SPREADER AND ACTIVATOR.

DIRECTIONS

1. Be sure soil is moist and lawn healthy before treating. Do NOT apply to newly seeded lawns or to lawns stunted and dormant because of drought. During drought, water grass well several days in advance, and restore turf to healthy green color with Comar Lawn Food before treatment.

2. Keep lawn mowed to about 1 1/2" to 2" during treatment.

HOW TO APPLY: Use hose gun, or watering can; coarse sprays are less likely to drift.

For Common Broad-Leaf Weeds, and Young Seeding Crabgrass use 1 tablespoon per gallon of water applied to 65 square feet of lawn at temperatures above 85 F., and 1 1/2 tablespoons per gallon of water per 65 square feet for temperatures below 85 F.

On older or branched crabgrass, or hard-to-kill weeds, use 2 tablespoons per gallon of water in hot weather, per 65 square feet of lawn. If weather is cool (70-80 F), double dosage in each case. The interval between treatments should be one week.

Lawn may slightly discolor from treatment, but if directions are followed, this should be only temporary, and your good turf grasses will quickly return to normal color after first cutting.

In cool weather, weeds may take some time to show effect, but do not be impatient, since kill will result. Do not apply until grass is dry. Apply on a still day, to avoid drift. Do not water until a day after application. Repeat treatment as needed, at 2 to 3 week intervals. Flush spray jar and head thoroughly after use with mild ammonia solution (2 tablespoons per gallon).

WILL NOT INJURE TURF WHEN USED AS DIRECTED

TO SPRAY LARGE AREAS: Measure out 1000 square feet (20' x 50') and place required number of tablespoons in hose sprayer jar. Fill jar to top with water, dissolving powder. Apply evenly to area, until jar is empty.

- KILLS BROADLEAF WEEDS FROM ROOT TO CROWN
- KILLS CRABGRASS