

Lot No.

1-21-72
6079-7

TOXAPHENE

EMULSIFIABLE CONCENTRATE

(Contains 6 lbs. Toxaphene Per Gallon)

DIRECTIONS FOR USE

This product emulsifies readily, forming a stable emulsion when added to water. Slight agitation will maintain a uniform dispersion.

COTTON INSECTS: In early season applications for thrips and flea hoppers apply 1½ to 3 pints per acre in 2 to 5 gallons of water and repeat at weekly intervals as long as infestations persist.

In pre-square applications for boll weevil control apply 2 to 3 quarts per acre in 2 to 5 gallons of water in 2 or 3 applications at weekly intervals. Consult State Agricultural Experiment Station regarding timing and adaptability of early season control programs in your locality.

For a mid season control program for boll weevil apply 2 to 3 quarts per acre in 2 to 5 gallons of water starting when up to 10% of the squares are punctured and repeat at 5 to 7 day intervals until 3 applications have been made or as long as weevil infestation persists.

ACTIVE INGREDIENTS:

Toxaphene Technical Chlorinated Camphene,
Chlorine Content 67-69

60.0%

Petroleum Distillate

36.0%

INERT INGREDIENTS:

4.0%

TOTAL

100.0%

WARNING

Keep this and all other containers tightly closed. Do not breathe dust. Avoid contact with skin and clothing. Do not get in eyes. If in eyes, flush with water. Do not eat, drink, or smoke while using this product. Wash thoroughly after use. Do not use in residential areas. Do not use near water. Do not use near children. Do not use near livestock. Do not use near poultry. Do not use near bees. Do not use near fish. Do not use near aquatic life. Do not use near birds. Do not use near mammals. Do not use near reptiles. Do not use near amphibians. Do not use near plants. Do not use near animals. Do not use near humans. Do not use near children. Do not use near pregnant women. Do not use near nursing infants. Do not use near the elderly. Do not use near the infirm. Do not use near the disabled. Do not use near the sick. Do not use near the dying. Do not use near the living. Do not use near the dead. Do not use near the buried. Do not use near the cremated. Do not use near the ashes. Do not use near the bones. Do not use near the teeth. Do not use near the hair. Do not use near the nails. Do not use near the skin. Do not use near the eyes. Do not use near the ears. Do not use near the nose. Do not use near the mouth. Do not use near the throat. Do not use near the lungs. Do not use near the stomach. Do not use near the intestines. Do not use near the bladder. Do not use near the uterus. Do not use near the vagina. Do not use near the penis. Do not use near the testicles. Do not use near the prostate. Do not use near the kidneys. Do not use near the liver. Do not use near the spleen. Do not use near the pancreas. Do not use near the gallbladder. Do not use near the stomach. Do not use near the intestines. Do not use near the bladder. Do not use near the uterus. Do not use near the vagina. Do not use near the penis. Do not use near the testicles. Do not use near the prostate. Do not use near the kidneys. Do not use near the liver. Do not use near the spleen. Do not use near the pancreas. Do not use near the gallbladder.

NET CONTENTS GALLONS
NORTH LOUISIANA CHEMICALS, INC.
MILLIAM, LOUISIANA

Lot No.

7-21-77
6074 7

TOXAPHENE

EMULSIFIABLE CONCENTRATE

(Contains 6 lbs. Toxaphene Per Gallon)

DIRECTIONS FOR USE

This product emulsifies readily when mixed with water. Add 1 quart of concentrate to 100 quarts of water. Repeat at weekly intervals as indicated in directions per pest.

COTTON INSECTS: In early season, apply 1 quart of concentrate and five quarts of water to 100 quarts of water. Repeat at weekly intervals as indicated in directions per pest.

In late season, apply 1 quart of concentrate and five quarts of water to 100 quarts of water in 2 or 3 applications at weekly intervals. Consult State Agricultural Experiment Station regarding timing and amount of application. Do not apply to cotton in bloom.

In mid season, apply 1 quart of concentrate and five quarts of water to 100 quarts of water in 2 or 3 applications at weekly intervals. Consult State Agricultural Experiment Station regarding timing and amount of application. Do not apply to cotton in bloom.

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphor)
Chlorine Content 67-69%
Petroleum Distillate

100.0

100.0

INERT INGREDIENTS:

3.0

TOTAL

100.0%

NET CONTENTS:

GALLONS

NORTH LOUISIANA CHEMICALS, INC.

GILLIAM, LOUISIANA

TOXAPHENE

EPA Reg. No. 6079-7

EMULSIFIABLE CONCENTRATE

(Contains 6 lbs. Toxaphene Per Gallon)

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene, Chlorine Content 67-69%)	60.0%
Petroleum Distillate	36.0%

INERT INGREDIENTS:

4.0%

TOTAL

100.0%

WARNING: Keep out of reach of children!

See Side Panel for ANTIDOTE and Additional WARNING STATEMENTS

WARNING: MAY BE FATAL IF SWALLOWED! Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after use. Avoid contamination of food and foodstuffs.

This product will kill fish and wildlife. Keep out of lakes, ponds and streams. Do not apply when rain is likely to occur. This product is toxic to bees and should not be applied when bees are actively visiting the area. Do not spray dry animals or animals being finished for slaughter or for export. Apply this product only as specified on the label.

Do not spray or apply to water, or to water that is being treated.

Do not apply to water that is being used for drinking, or to water that is being used for irrigation. Do not apply to water that is being used for the purpose of raising shellfish. Do not apply to water that is being used for the purpose of raising fish. Do not apply to water that is being used for the purpose of raising aquatic plants. Do not apply to water that is being used for the purpose of raising aquatic animals. Do not apply to water that is being used for the purpose of raising aquatic insects. Do not apply to water that is being used for the purpose of raising aquatic mollusks. Do not apply to water that is being used for the purpose of raising aquatic arthropods. Do not apply to water that is being used for the purpose of raising aquatic nematodes. Do not apply to water that is being used for the purpose of raising aquatic rotifers. Do not apply to water that is being used for the purpose of raising aquatic protozoans. Do not apply to water that is being used for the purpose of raising aquatic fungi. Do not apply to water that is being used for the purpose of raising aquatic bacteria. Do not apply to water that is being used for the purpose of raising aquatic viruses. Do not apply to water that is being used for the purpose of raising aquatic parasites. Do not apply to water that is being used for the purpose of raising aquatic predators. Do not apply to water that is being used for the purpose of raising aquatic prey. Do not apply to water that is being used for the purpose of raising aquatic competitors. Do not apply to water that is being used for the purpose of raising aquatic symbionts. Do not apply to water that is being used for the purpose of raising aquatic mutualists. Do not apply to water that is being used for the purpose of raising aquatic commensals. Do not apply to water that is being used for the purpose of raising aquatic parasites. Do not apply to water that is being used for the purpose of raising aquatic predators. Do not apply to water that is being used for the purpose of raising aquatic prey. Do not apply to water that is being used for the purpose of raising aquatic competitors. Do not apply to water that is being used for the purpose of raising aquatic symbionts. Do not apply to water that is being used for the purpose of raising aquatic mutualists. Do not apply to water that is being used for the purpose of raising aquatic commensals.

NET CONTENTS:

GALLONS

NORTH LOUISIANA CHEMICALS, INC.

GILLIAM, LOUISIANA

DIRECTIONS FOR USE (Con't.)

For late season control of boll weevil apply 1/2 to 2 3 gallons per acre in 2 to 5 gallons of water, making sure adequate coverage is obtained depending on size of cotton plant. Repeat applications at 5 to 7 day intervals as long as infestation warrants.

North Louisiana Chemicals, Inc. warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this product is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

WARNING: Keep out of reach of children!

ANTIDOTE: Internal - Give one to two tablespoons of salt in glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

EXTERNAL - Wash with soap and water.

FLAMMABLE MIXTURE: Keep away from heat or flame.