

# EIGHT - TWO TOXAPHENE - METHYL PARATHION

NO. 67911

## EMULSIFIABLE CONCENTRATE

Contains 8.2 lbs. Parathion Per Gallon

### ACTIVE INGREDIENTS

Toxaphene	66.6
Methyl Parathion	16.8
Emulsifier	16.6

### INERT INGREDIENTS

Total	100
-------	-----

North Louisiana Chemicals, Inc. warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this product is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

NET CONTENTS                      GALLONS

### NORTH LOUISIANA CHEMICALS, INC.

MONROE, LOUISIANA

DIR

DATE

S

# EIGHT - TWO

# TOXAPHENE - METHYL PARATHION



POISON

EPA REG. NO. 60-407

LOT NO.

RATE

Per Gallon.

## DIRECTIONS FOR USE

this product can be used in conjunction with other pesticides. When used in conjunction with other pesticides, this product should be used in accordance with the directions for use of the other pesticides. The use of this product in conjunction with other pesticides should be made only when the use of this product is necessary to control the pest being treated. The use of this product in conjunction with other pesticides should be made only when the use of this product is necessary to control the pest being treated.

Directions for use text, including application rates and safety instructions. The text is mostly illegible due to low contrast and scan quality. Key phrases like 'Apply to the foliage of the plant' and 'Workers should wear protective clothing' are faintly visible.

S INC.

# EIGHT - TWO

# TOXAPHENE - METHYL PARATHION

ITRA 1-

DIRTY 12 1957

Directions: For Use: Or Cotton

Apply to the undersides of leaves and stems of cotton plants.

Apply during the early stages of growth and repeat applications at 7 to 10 day intervals. Do not apply to cotton plants during the flowering stage.

Mid Season: Apply to cotton plants during the flowering stage. Do not apply to cotton plants during the fruiting stage.

Pre-Season: Apply to cotton plants during the pre-season stage. Do not apply to cotton plants during the flowering stage.

Apply to cotton plants during the flowering stage. Do not apply to cotton plants during the fruiting stage.

Apply to cotton plants during the flowering stage. Do not apply to cotton plants during the fruiting stage.