USE DIRECTIONS

FOR CLEANING

Lightly soiled areas — surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of Germicidal Cleaner per gallon of water. Rinse cleaned area with fresh water.

Heavily soiled areas — mop or brush soiled surface with a solution containing 2 ounces of Germicidal Cleaner per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing 1 ounce of Germicidal Cleaner per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces of Germicidal Cleaner per gallon of water and then disinfected. All treated food contact surfaces and/or equipment must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse. To disinfect locker rooms, shower stalls, walkways around pools and other areas against pathogenic fungi, mop area with a solution containing 1 ounce of Germicidal Cleaner in 1 gallon of water. This solution will destroy Trichophyton mentagrophytes, a causative agent of athlete's foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 1 ounce of Germicidal Cleaner per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

GERMICIDAL CLEANER

Phenol Coeff: S. aureus - 15

S. typhosa — 9

DISINFECTANT-CLEANER-SANITIZER-FUNGICIDE-DEODORANT

ACTIVE INGREDIENTS:

Isopropyl Alcohol	10.00%
sodium o-phenylphenate	
sodium 4-chloro-2-cyclopentylphenate	3.89%
sodium dodecylbenzene sulfonate	2.90%
tetrasodium ethylene diamine tetraacetate	0.20%

INERT INGREDIENTS 79.06%

EPA Reg. No. 6125-12

DANGER

KEEP OUT OF REACH OF CHILDREN

See First Aid statement and additional precautions on side panel. Container Disposal: When empty, rinse container thoroughly with water and discard in a safe place.

DO NOT USE, POUR, SPILL OR STORE NEAR OPEN FLAME.

SANITIZATION

To sanitize wall, floors and a food processing plants, first oughly with a solution contain of Germicidal Cleaner per gal If surface is heavily soiled, us Cleaner at a rate of 2 ounces insure thorough cleaning. San surface with a rinse solution ounce of Germicidal Cleaner water. All treated food contand/or equipment must be scrubbed with detergents and potable water prior to reuse.

DEODORIZATION

To deodorize surfaces during of or brush surfaces with a solu ing 2 ounces of Germicidal gallon of water. Germicidal C effective antibacterial cleaner use in household, industrial tional cleaning. When used Germicidal Cleaner will disinfe faces, such as floors, walls, and furniture. It may be rel deodorize bathroom areas, ga food storage areas, kennel are areas where obnoxious odors This product may be used a purpose antibacterial-antifung in homes, schools, restaurant nursing homes, public rooms, ing plants and in other place cient cleaning and antimicrot

See left panel for use direction

DANGER

desired.

KEEP OUT OF REACH OF

Corrosive. Causes eye damagirritation. Do not get in eyes on clothing. Protect eyes and handling. Harmful if swallowed tamination of food.

FIRST AID

In case of contact, immediate or skin with plenty of water 15 minutes. For eyes, call Remove and wash contaminate before reuse.

If swallowed, drink promptly a tity of milk, egg whites, gelati

DNS

s — surfaces to be cleanpped or brushed with a g 1 ounce of Germicidal of water. Rinse cleaned later.

is — mop or brush soiled ition containing 2 ounces iner per gallon of water. face with fresh water.

DSE DISINFECTION

floors and other hard ls, schools, restaurants, her public areas, mop or e treated with a solution e of Germicidal Cleaner ter. Surfaces that are buld be cleaned with 2 cidal Cleaner per gallon disinfected. All treated laces and/or equipment ly scrubbed with deterwith potable water prior ect locker rooms, shower around pools and other hogenic fungi, mop area containing 1 ounce of er in 1 gallon of water. Il destroy Trichophyton causative agent of ath-

NFECTION

ls, floors, furniture and es in hospitals and nursor brush surface to be ution containing 1 ounce aner per gallon of water, at are heavily soiled with a rial or exudates should brubbed with a solution lees per gallon of water with a fresh solution conter gallon of water before

GERMICIDAL CLEANER

Phenol Coeff: S. aureus - 15

S. typhosa - 9

DISINFECTANT-CLEANER-SANITIZER-FUNGICIDE-DEODORANT

ACTIVE INGREDIENTS:

Isopropyl Alcohol	10.00%
sodium o-phenylphenate	
sodium 4-chloro-2-cyclopentylphenate	3.89%
sodium dodecylbenzene sulfonate	2.90%
tetrasodium ethylene diamine tetraacetate	0.20%
INERT INGREDIENTS	79.06%

EPA Reg. No. 6125-12

DANGER

KEEP OUT OF RLACH OF CHILDREN

See First Aid statement and additional precautions on side panel. Container Disposal: When empty, rinse container thoroughly with water and discard in a safe place.

DO NOT USE, POUR, SPILL OR STORE NEAR OPEN FLAME.

SANITIZATION

To sanitize wall, floors and equipment in food processing plants, first clean thoroughly with a solution containing 1 ounce of Germicidal Cleaner per gallon of water. If surface is heavily soiled, use Germicidal Cleaner at a rate of 2 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing 1 ounce of Germicidal Cleaner per gallon of water. All treated food contact surfaces and/or equipment must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.

DEODORIZATION

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of Germicidal Cleaner per gallon of water. Germicidal Cleaner is an effective antibacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, Germicidal Cleaner will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnexious odors may develop. This product may be used as a general purpose antibacterial-antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

See left panel for use directions.

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or

with fresh water.

Phenol Coeff: S. aureus - 15

S. typhosa - 9

DISINFECTION

oors and other hard schools, restaurants, public areas, mop or eated with a solution f Germicidal Cleaner Surfaces that are be cleaned with 2 Cleaner per gallon infected. All treated and/or equipment crubbed with deterpotable water prior locker rooms, shower nd pools and other nic fungi, mop area taining 1 ounce of 1 gallon of water. estroy Trichophyton usative agent of ath-

CTION

loors, furniture and hospitals and nurstrush surface to be containing 1 ounce per gallon of water. The heavily soiled with or exudates should bed with a solution per gallon of water a fresh solution contallon of water before

DISINFECTANT-CLEANER-SANITIZER-FUNGICIDE-DEODORANT

ACTIVE INGREDIENTS:

Isopropyl Alcohol	
sodium o-phenylphenate	
sodium 4-chloro-2-cyclopentylphenate	3.89%
sodium dodecylbenzene sulfonate	2.90%
tetrasodium ethylene diamine tetraacetate	0.20%
INERT INGREDIENTS	79.06%


EP# Reg. No. 6125-12

DANGER KEEP OUT OF REACH OF CHILDREN

See First Aid statement and additional precautions on side panel. Container Disposal: When empty, rinse container thoroughly with water and discard in a safe place.

DO NOT USE, POUR, SPILL OR STORE NEAR OPEN FLAME.

MANUFACTURED BY BIXON CHEMICAL COMPANY 50-19 97th Place Corona, New York 11368



NOVO 2 1973

FUNGICIDE ALL ENTICIDE ACT FOR ECONOLIC POISON REGISTERED UNDER NO. 6/25-12

potable water prior to reuse.

DECDORIZATION

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of Germicidal Cleaner per gallon of water. Germicidal Cleaner is an effective antibacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, Germicidal Cleaner will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial-antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

See left panel for use directions.

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.