

[6125-16]

DIRECTIONS FOR USE INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly before using.

Vacate treated area and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

Surface Spraying: To control Roaches, Ants, Silver Fish, Spiders, Crickets, Clover Mites, Cheese Mites and exposed stages of Granary Weevils, and exposed stages of Rice Weevils, and exposed stages of Flour Beetles, and exposed stages of Rust Red Flour Beetles, and exposed stages of Saw Toothed Grain Beetles, and exposed stages of Spider Beetles, and exposed stages of Cigarette Beetles, and exposed stages of Drugstore Beetles, and exposed stages of Meal Worms, and exposed stages of Grain Mites, and exposed stages of Caddisflies. Use a good spray adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under panels, around containers of stored food, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases, for ants, spray trails, nests and points of entry. At surface application areas to be used, spray floors, walls, and other surface applying at a rate of 1 gallon to 750 square feet of surface.

To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks and infested areas of shelving. Repeat treatment as needed.

To control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed.

To control Fleas and Brown Dog Ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

Jan. 2, 1975

PYRENONE[®]

General Purpose Aqueous Insecticide

EPA Reg. No. 6125-16

A Concentrated PYRENONE[®] Micro Emulsion

— Clear

— Stable

— Leaves No Oily or Objectionable Residue

Effective, Fast Acting Insecticide.

For Use in Homes, Institutions, Restaurants.

Food Processing Plants, Dairies, Livestock and Outdoors
for Mosquito Control.

Apply with Conventional Mechanical or Compressed Air
Equipment.

ACTIVE INGREDIENTS:

Pyrethrins	1.1%
Piperonyl Butoxide, technical	1.0%
Petroleum distillate	4.4%

INERT INGREDIENTS:

Equivalent to 0.8% (butylcarbityl) (6-pronyl piperonyl) ether and to 0.7% related compounds

Pyrenone is a registered trademark of FMC Corporation

CAUTION

Keep Out Of Reach Of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

MANUFACTURED BY
BIXON CHEMICAL COMPANY

50-19 97th PLACE

CORONA, NEW YORK 11368

Net Contents Gallons

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining. Special attention should be given to seams and folds.

Space Spraying: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of Flies, Fruit Flies, Gnats, Mosquitoes, Wasps, Hornets, and exposed stages of Angoumois Grain Moths, and exposed stages of Mediterranean Flour Moths, and exposed stages of Indian Meal Moths, and exposed stages of Tobacco Moths: direct spray at an upward angle, distributing it uniformly throughout the entire area, at a rate of 3-6 ounces per 1000 cubic feet of space.

Outdoor Adult Mosquito Control: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited. Apply at a rate of 1½ gallons per acre (4½ ounces per 1000 square feet).

Using truck mounted conventional misting equipment such as Rotomist Models 51C, 91 or 100 ht. traveling at 5 m.p.h. to treat a 300 foot swath, calibrate equipment to deliver 4.5 gallons per minute.

CAUTION: Harmful if swallowed. Avoid contamination of food and foodstuffs. Remove pets, birds and cover fish aquaria before spraying.

This product is toxic to fish. Keep out of lakes, streams or ponds.

Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.