

**Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations.**

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard surfaces. No potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. Quat No. 8 in 4 gallons of water.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush. No potable water rinse is required.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz./4 gallon dilution of Quat No. 8. No Potable water rinse is required.

At 1 oz./4 gallons, Quat No. 8 fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinances 1965 Recommendations of the U.S. Public Health Services in waters up to 1000 ppm of hardness calculated as Ca CO<sub>3</sub> when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. Quat No. 8 in 4 gallons of warm water. No Potable water rinse is required.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER**

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

**STORAGE AND DISPOSAL**

- DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

**PESTICIDE DISPOSAL**

PESTICIDE OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

**CONTAINER DISPOSAL**

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS, OR BURY IN A SAFE PLACE.

**GENERAL**

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

**QUAT No. 8  
DISINFECTANT - SANITIZER  
FUNGICIDE DEODORIZER**

Disinfectant-Sanitizer-Fungicide  
Deodorizer for Hospital, Institutional,  
Industrial, School, Dairy and Other  
Farm and Home Use

**AOAC Phenol Coefficients**


Staphylococcus aureus, ATCC #6538	77.5
Salmonella typhosa, ATCC #6539	77.5

**Active Ingredients**

Didecyl dimethyl ammonium chloride	7.5%
Isopropanol	3.0%

**Inert Ingredients**

	89.5%
	100.0%



**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

**Statement of Practical Treatment**

In case of contact, immediately flush eyes of skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

**SEE LEFT PANEL FOR**

**ADDITIONAL PRECAUTIONARY STATEMENTS**

E.P.A. Est. #6125 N.Y. 01  
E.P.A. Registration #6125-31  
Net Contents

Manufactured By:  
**BIXON CHEMICAL CO.**  
50-19 97th Place  
Corona, N.Y. 11368

**Directions for Use**

**GENERAL CLASSIFICATION**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

006

**Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions.**

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces.

Add 3 1/2 oz. Quat No. 8 to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use-level, Quat No. 8 is effective against Pseudomonas aeruginosa.

**Disinfectant in Institutions, Industry, Schools and Homes**

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

Add 2 oz. of Quat No. 8 to 4 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 2 oz./4 gallon use-level, Quat No. 8 is effective against Staphylococcus aureus, Salmonella choleraesuis and Trichophyton interdigitale (the athlete's foot fungus).

**Disinfection of Barber Tools**

Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2-oz./gallon solution of Quat No. 8.

**Disinfection of Poultry Equipment, Animal Quarters and Kennels.**

Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. Quat No. 8 in 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

60

60A