

BEST DOCUMENT AVAILABLE

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

**FIRST AID:** If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

**STORAGE & DISPOSAL:** Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Residue that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water sources. Open dumping is prohibited.

**ALGAECIDE N.F. 6**

**ACTIVE INGREDIENT**

Poly(oxyethylene (dimethylamino) ethylene-  
(dimethylamino) ethylene dichloride) ..... 0.0%

**INERT INGREDIENT** ..... 94.0%

This product contains 0.5 lb. of active ingredient per gallon and weighs 6.44 lb. per gallon.

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

EPA REG. NO. 6125-42  
EPA EST. NO. 6125 N.Y. 01  
NET CONTENTS

Manufactured by  
**CHEMICAL COMPANY**  
30-18 57th Place  
Corona, N.Y. 11368

ACCEPTED  
6125-42  
FEB 22 1980  
UNDER THE FEDERAL INSECTICIDE  
HERBICIDE AND RODENTICIDE ACT  
FOR RECORDING PURPOSES  
ED 100818-20

**DIRECTIONS**

It is a solution of a water-insoluble active ingredient. ALGAECIDE N.F. 6 is a water-soluble concentrate. It is used to control algae in lakes, streams, ponds, and other bodies of water. It is also used to control algae in industrial water systems. It is a contact herbicide. It is not a systemic herbicide. It is not a soil herbicide. It is not a foliar herbicide. It is not a root herbicide. It is not a stem herbicide. It is not a leaf herbicide. It is not a fruit herbicide. It is not a seed herbicide. It is not a soil sterilant. It is not a soil fumigant. It is not a soil disinfectant. It is not a soil sanitizer. It is not a soil conditioner. It is not a soil amendment. It is not a soil improver. It is not a soil enhancer. It is not a soil builder. It is not a soil stabilizer. It is not a soil protector. It is not a soil preserver. It is not a soil restorer. It is not a soil rejuvenator. It is not a soil revitalizer. It is not a soil reclaimer. It is not a soil regenerator. It is not a soil renewer. It is not a soil restorer. It is not a soil rejuvenator. It is not a soil revitalizer. It is not a soil reclaimer. It is not a soil regenerator. It is not a soil renewer.

BEST DOC

**DIRTY**  
**GENERAL**  
In violation of Federal law to use this product in a manner inconsistent with its labeling.  
**ALGAEICIDE N.F. 6**  
Algae, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove sludge, growth, microbiological films, and other deposits. An initial slug addition of 10 to 20 fluid ounces of ALGAEICIDE N.F. 6 per 1000 gallons of water to provide a concentration of 80 to 200 parts per million of ALGAEICIDE N.F. 6, based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.  
Subsequent slug additions of 2.50 to 25.00 fluid ounces of ALGAEICIDE N.F. 6 per 1000 gallons of water (20 to 200 parts per million of ALGAEICIDE N.F. 6) should be employed every 3 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.  
ALGAEICIDE N.F. 6 is used to control bacteria in industrial air-washing systems that maintain effective mist-eliminating components. Prior to its use, systems should be cleaned to remove bacterial slime and other deposits. An initial slug dose of 37.12 to 62.44 fluid ounces of ALGAEICIDE N.F. 6 per 1000 gallons of water is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 20.31 to 62.44 fluid ounces of ALGAEICIDE N.F. 6 per 1000 gallons of water should be employed each 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and severity of the bacterial problem. Slug additions may be made to the sump or to the water collection trays of the airwash system.

**PRECAUTIONS**  
Avoid breathing dust or fumes. Avoid contact with skin, eyes, or clothing.  
If swallowed, drink promptly a large glass of water, or milk, or other liquid. If it is in the eyes, flush with large quantities of water. If on skin, wash immediately.  
**HAZARDS:** This pesticide is toxic to fish, birds, or ponds. Permit no discharges containing this material into streams, ponds, or public water. For the regional office of the EPA, contact the nearest office of the EPA.

BEST DOCUMENT AVAILABLE

# ALGAEICIDE N.F. 6

**ACTIVE INGREDIENT**  
Poly(oxyethylene dimethyliminio) ethylene-  
(dimethyliminio) ethylene dichloride ..... 6.0%

**INERT INGREDIENT** ..... 94.0%

This product contains 0.5 lb. of active ingredient per gallon and weighs 8.44 lb. per gallon

## CAUTION

KEEP OUT OF REACH OF CHILDREN

EPA REG. NO. 6125-40  
EPA EST. NO. 6125 N.Y. 01  
NET CONTENTS

Manufactured by  
**CHEMICAL COMPANY**  
50-19 97th Place  
Corona, N.Y. 11368

**ACCEPTED**  
**6725-40**  
**FEB 22 1980**  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ACCURATE WEIGHT MEASUREMENT  
BY NIBBS, INC.

### DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
ALGAEICIDE N.F. 6 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove sludge, growth, microbiological films, and other deposits. An initial slug addition of 10 to 20 fluid ounces of ALGAEICIDE N.F. 6 per 1000 gallons of water to provide a concentration of 80 to 200 parts per million of ALGAEICIDE N.F. 6, based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.  
Subsequent slug additions of 2.50 to 25.00 fluid ounces of ALGAEICIDE N.F. 6 per 1000 gallons of water (20 to 200 parts per million of ALGAEICIDE N.F. 6) should be employed every 3 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.  
ALGAEICIDE N.F. 6 is used to control bacteria in industrial air-washing systems that maintain effective mist-eliminating components. Prior to its use, systems should be cleaned to remove bacterial slime and other deposits. An initial slug dose of 37.12 to 62.44 fluid ounces of ALGAEICIDE N.F. 6 per 1000 gallons of water is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 20.31 to 62.44 fluid ounces of ALGAEICIDE N.F. 6 per 1000 gallons of water should be employed each 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and severity of the bacterial problem. Slug additions may be made to the sump or to the water collection trays of the airwash system.

BEST DOCUMENT AVAILABLE