
LOFTS CHINCH KILLER with Diazinon SCOTT CYCLONE **KILLS** SEARS CHINCH BUGS CLOVER MITES SOD WEBWORMS (LAWN MOTHS) JAPANESE BEETLE GRUBS

ACTIVE INGREDIENT *O,O-diethyl O-(2-isopropyl-6 methyl- 4pyrimidinyl) phosphorothioate **INERT INGREDIENTS:**

U. 5 Partie From 2.754 233 *Diazinon * Trademark of CIBA-GE/GT Corp.

DIRECTIONS - APPLY WITH SPREADER



1.50% 98.50%

FISH AND WILDLIFE CAUTION

1972

This product is toxic to fish and wildlife. Birds

ACTIVE INGREDIENT

*O,O-diethyl O-(2-isopropyl-6 methyl- 4

pyrimidinyl) phosphorothioate

INERT INGREDIENTS:

•

U. S. Patent No. 2,754,243 *Diazinon * Trademark of CIBA-GE'GY Corp

DIRECTIONS --- APPLY WITH SPREADER

LAWNS: For control of CHINCH BUGS apply at the rate of 15 pounds per 2500 square feet; as soon as the pest appears. Repeat in 7 to 10 days if necessary.

Granular Chinch Bug Killer applied at the above rate also efficiently controls Sod Webworms, ants. fleas, chiaaers and CLOVER MITES that infest lawns.

In the New York Metropolitan area, Chinch Bugs have one generation in June and a second in August. Consult your Extension Service Specialist as to exact timing of applications.

For control of JAPANESE BEETLE GRUBS, EUROPEAN CHAFER GRUBS, and SOUTHERN CHAFER GRUBS apply at the rate of 15 pounds per 1685 square feet.

For Grub control, apply anytime between late July and early October. For control of HYPERODES WEEVIL (New York State only), apply at the rate of 15 pounds per 1850 square feet. Treat problem areas in mid April and again in mid May.

NOTE: After application sprinkle the lawn thoroughly to wash the insecticide off the grass and on to the turf. Keep children and pets off treated area until lawn has been watered and completely dried.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid breathing dust. Harmful if swallowed, May be absorbed through skin. Avoid storage near food and thed products. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling:

Do not the empty container. Destroy by 1 Hying with wastes or burning. ' y away from smoke or fumer.

1.50% **98.50%**

FISH AND WILDLIFE CAUTION

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

EPA Reg No. 6143

Manufactured For

PEDIGREED LOFTS SEED COMPANY

BOUND BROOK, NEW JERSEY

NET WEIGHT/5 POUNDS

· · · ·