LUFTIS

DANDELICH AND BROADLEAF

LAWN WED KILLER

Such as: Carpellons, Plantain, Bucharn, Chickeed, Knoweed, Surrel and many others.

Ingredient Statement

HCRBICIDAL ACTIVE INCREDIENTS Dimethylamine Salts of مر 2،4.4 2,4-Dichlorophenoxyacetic Acid* Isooctyl Ester of Silvex (2-(2,4,5-Trichloropnenuxy) Propionic Acid)** م 1.32 م INERT INGREDIENTS (Other than Herbicidal) 94.06/2 مُر ١٥٥٠٥٥ مُر TOTAL

* Equivalent to 3.84% 2,4-Dichlorophenoxyacetic Acid

** Equivalent to .94, 3ilvex

U.3.D.A. REG. #_

COVERS 5,000 **S**QUARE FEET (501 x 1001) CAUTION: Keep out of Reach of children. See back Panel for Additional

Cautions.

NET WEIGHT 6 POUNDS

<u>uASK</u>SE uAu

NET METURE & PULLIUS

LOF ITS

DANDELIUM AND EMCALETAN

LAWN WEED KILLER

<u>Ingredient</u> statement

HEREICIDAL

ACTIVE INCREDIENTS

Dimethylamine Salts of

2, 4-Dichlorophenoxyacetic Acid * 4.62%

Isooctyl Ester of Silvex

(2-(2,4,5-Trichlorophenoxy) Propionic Acid) ** 1.32%

INERT INCREDIENTS (Other than Herbicidal) 94.06%

TOTAL 100.00%

*Equivalent to 3.84, 2,4-Dichlorophenoxyacetic Acid ** Equivalent to .94, Silvex

> LOFT'S PEDIGREED SEED COMPANY CHINNEY ROCK ROAD BOUND BROOK, NEW JERSEY

Directions for use

2,4-D Aming and Silvex on a lightweight carrier. Its function is to control broadleaf weeds such as:

Plantain Danvelion Wild Carrot
Buckhorn Common Chickweed Wild Onion
Thistle Mouse cared Chickweed Pigweed
Rappeed Sou Grass Lambs Quarters
White Clover Clover And many other

The combination of ingredients is very effective, economical and easy to use.

LOFT'S CANCELION AND BROADLEAF LAW'S WELD KILLER can be used in the Spring, Summer of Fall. However, the best results are obtained when the weeds are growing vigorously. When using this product strive to distribute it uniformly and to retain the granules on the leaves of weeds. The following suggestions will assist:

- 1. The contents of this way are to cover a 5000 sq. ft. area $(*0* \times 100*)$. This can best be accomplished using a spreader.
- 2. Ap to between mowings to provide sufficient leaf surface to be pated.
- 3. Sprinkle lawn prior to application or apply after a light rainful.
- 4. Lela, application when rain is predicted.
- 5. Do not water lawn for 24 hours after application.
- 6. Avoid application on windy days to avoid drift to surrounding plants.
- 7. Do not treat newly seeded lawns until grasses are well established or at least until after third mowing.

SUGGESTED SPREADER SETTINGS

Lawn Seauty	8
Scott*s	6
Jackson	5
Cyclone	3

These suggestions are for your guidance in spreading, and are in no sense a guarantee of exact application. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in the rates applied.

CAUTION: Avoid innalation or contamination of food stuffs. Avoid contact with skin, eyes or clotning. Wash hands after using. To not some near fertilizer, seeds, insecticides or fungicides. Do not allow dust to drift onto susceptible plants such as ornamentals, vegetables, etc. Do not apply to stoloniferous grasses (those with underground stems) such as bents, bermuda or St. Augustine. Do not use on clover if you desire it a remain in the lawn. Certain desirable grasses may exhibit slight browning but will recover after a short period if the product has been used as directed. Vapors from this product may cause injury to susceptible plants growing in the immediate vicinity.

BACK OF

NET WEIGHT 6 POU

LOFT'S

DANDELION AND B

LAWN WEED KI

Ingredient Statem

HERBICIDAL

Dimethylamine Salts of

2, 4-Dichlorophenoxyacetic
Isooctyl Ester of Silvex
(2-(2,4,5-Trichlorophenox
INERT INGREDIENTS (Other tha

*Equivalent to 3.84% 2,4-Dichlorop
** Equivalent to .94% Silvex

LOFT'S PEDIGREED S CHINNEY ROCK BOUND BROOK, N

Directions

2,4-D Amine and Silvex on a lightwell control broadleaf weeds such as:

Plantain
Buckhorn
Thistle
Ragweed
White Clover

Dand Comm Mous Sour Clov

The combination of ingredients is verto use.

LOFT'S DANDELION AND BROADLEAF LAWN Summer of Fall. However, the best rare growing vigorously. When using uniformly and to retain the granules suggestions will assist:

- 1. The contents of this bag are to cover a 5000 sq. ft. area (50° x 100°). This can best be accomplished using a spr2ader.
- 2. Apply between mowings to provide sufficient leaf surface to be treated.
- 3. Sprinkle lawn prior to application or apply after a light rainfall.
- 4. Delay application when rain is predicted.
- 5. Do not water lawn for 24 hours after application.
- 6. Avoid application on windy days to avoid drift to surrounding plants.
- 7. Do not treat newly seeded lawns until grasses are well established or at least until after third mowing.

SUGGESTED SPREADER	SETTINGS
Lawn Beauty	8
Scott*s	6
Jackson	5
Cyclone	3

These suggestions are for your guidance in spreading, and are in no sense a guarantee of exact application. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in the rates applied.

CAUTION: Avoid inhalation or contamination of food stuffs. Avoid contact with skin, eyes or clothing. Wash hands after using. Bo not store near fertilizer, seeds, insecticides or fungicides. Bo not allow dust to drift onto susceptible plants such as ornamentals, vegetables, etc. Do not apply to stoloniferous grasses (those with underground stems) such as bents, bermuda or St. Augustine. Do not use on clover if you desire it to remain in the lawn. Certain desirable grasses may exhibit slight browning but will recover after a short period if the product has been used as directed. Vapors from this product may cause injury to susceptible plants growing in the immediate vicinity.