

LOFTS

PROFESSIONAL

CRABGRASS

AND POA ANNUA

KILLER

WITH BALAN™

ACCEPTED
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
UNDER NO

A selective pre-emergence herbicide for the control of crabgrass, annual bluegrass (poa annua), goosegrass, and other annual weed grasses in established:

Bluegrass (perennial)

Ryegrass (perennial)

Fescue

Bahagrass

Bermudagrass

Centipedegrass

St. Augustinegrass

Zoysiagrass

TEST
SITE

THIS PRODUCT IS ON A CORN COB CARRIER-CAN BE USED IN A ROTARY TYPE (CYCLONE) OR DROP-TYPE SPREADER.

WITH BALANCE

REGISTERED
IN EQUIN...
UNDE...

...
...
...

Buggress, perennials
Ryegrass, perennials
Fescue
Barngrass


Perennial ryegrass
Candytuft
St. Augustine grass
Zoysia grass

THIS PRODUCT IS ON A CORN COB CARRIER-CAN BE USED IN A ROTARY TYPE
(CYCLONE) OR DROP-TYPE SPREADER.

NET WEIGHT 20 POUNDS

TREATS 5,000 SQ. FT.

CAUTION
Keep Out Of Reach Of Children.
See Back Panel For Additional Cautions



ACTIVE INGREDIENT	
N-butyl-N-ethyl-2,2,2-Trifluoro-2,6-dinitro-p-toluidine	1.15%
INERT INGREDIENTS (relative to herbicidal content)	98.85%
	<u>100.00%</u>

EPA Reg. No. 6143-19 EPA Est. 1159-NJ-1

BALANCE is the trade mark for Super — ELANCO PRODUCTS COMPANY
A DIVISION OF SQUIBB & COMPANY

NET WEIGHT 20 POUNDS

LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER WITH BALAN™.

A selective pre-emergence herbicide for the control of crabgrass, annual bluegrass (poa annua), goosegrass, and other annual weed grasses in established:

BLUEGRASS (perennial)
RYEGRASS (perennial)
FESCUE
BAHIAGRASS

BERMUDAGRASS
CENTIPEDEGRASS
ST. AUGUSTINEGRASS
ZOYSIAGRASS

ACTIVE INGREDIENT

N-butyl-N-ethyl-a,a,a-Trifluoro-2,6-dinitro-p-toluidine ----- 1.15%

INERT INGREDIENTS (relative to herbicidal content) ----- 98.85%
100.00%

EPA Reg. No. 6143-19 EPA Est. 1159-NJ-1

BALAN™ — the trade mark for Benefin — ELANCO PRODUCTS COMPANY
A DIVISION OF ELI LILLY & COMPANY

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with

GENERAL INFORMATION

LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER has the following advantages:

1. **CONVENIENT & ECONOMICAL** — LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER is on a corn cob carrier — can be used effectively in a rotary type spreader as well as a drop type spreader.
2. **GOOD SAFETY MARGIN** — Applied as directed, will not injure perennial Bluegrass, perennial Ryegrass, Fescue, Bermudagrass, Zoysiagrass, St. Augustinegrass, Centipede and Bahiagrass. Do not apply to new lawns started the previous fall.
3. **ESTABLISHES A WEED CONTROL ZONE** — LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER establishes a weed control zone by killing certain annual weed seeds as they germinate.
4. **WEEDS CONTROLLED** — Application prior to weed germination will prevent smooth and hairy crabgrass, annual bluegrass (poa annua), goosegrass (silver crabgrass or crowfoot), watergrass (barnyardgrass) and yellow and green foxtail.

WHEN TO APPLY:

FOR CRABGRASS CONTROL — Apply LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER in the spring before crabgrass germinates. Reseeding should be delayed for 90 to 120 days after application. To insure satisfactory reseeding results after using a pre-emergence chemical, the amount of seed should be increased.

FOR POA ANNUA (ANNUAL BLUEGRASS) CONTROL — Apply LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER in late to early fall (mid-July to mid-September) prior to annual bluegrass (poa annua) germination. Chemicals that effectively control poa annua seeds may thin established poa annua and fine-leaved fescues. Reseeding should be delayed 12 to 16 weeks after application.

HOW TO APPLY:

Apply LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER evenly over the turf area without skips or misses. A more uniform application can be made by spreading half the required amount of this product over the area, and then applying the remaining half at right angles or crosswise. Both drop-type and rotary-type spreaders may be used to apply LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER.

annual weed grasses in establishment:

**BLUEGRASS (perennial)
RYEGRASS (perennial)
FESCUE
BAHIAGRASS**

**BERMUDAGRASS
CENTIPEDEGRASS
ST. AUGUSTINEGRASS
ZOYSIAGRASS**

ACTIVE INGREDIENT

N-butyl-N-ethyl-a,a-Trifluoro-2,6-dinitro-p-toluidine ----- 1.15%

INERT INGREDIENTS (relative to herbicidal content) ----- 98.85%

100.00%

EPA Reg. No. 6143-19 EPA Est. 1159-NJ-1

BALANTM — the trade mark for Beneth — ELANCO PRODUCTS COMPANY
A DIVISION OF ELI LILLY & COMPANY

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with water. Do not contaminate foodstuffs or feed.


FISH AND WILDLIFE CAUTION

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes.

Do not reuse empty container. Destroy by burying with wastes or burning. Stay away from smoke or fumes.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

DESTROY BAG WHEN EMPTY. DO NOT REUSE.



PROFESSIONAL CRABGRASS AND POA ANNUA KILLER in late to early fall (mid-July to mid-September) prior to annual bluegrass (poa annua) germination. Chemicals that effectively control poa annua seeds may thin established poa annua and fine-leaved fescues. Reseeding should be delayed 12 to 16 weeks after application.

HOW TO APPLY:


Apply LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER evenly over the turf area without skips or misses. A more uniform application can be made by spreading half the required amount of this product over the area, and then applying the remaining half at right angles or crosswise. Both drop-type and rotary-type spreaders may be used to apply LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER.

Listed below are several suggested spreader settings. These should serve only as initial starting points for calibration since all spreaders will vary in their operation depending on the age, condition, particular model, and speed of operation.

APPROXIMATE SPREADER SETTINGS FOR LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER
(for applying the contents of this bag in one pass over the turf)

SUGGESTED SPREADER SETTINGS

- CYCLONE ----- 5
- O. M. SCOTT ----- 4
- LAWN BEAUTY ----- 5½



annual weed grasses in established:

BLUEGRASS (perennial)
 RYEGRASS (perennial)
 BROME
 FESCUE

BERMUDAGRASS
 CENTIPEDEGRASS
 ST. AUGUSTINEGRASS
 COYSIAGRASS

.....	1.15%
.....	98.85%
.....	100.00%

EPA Reg. No. 4102-10 EPA Est. 1159-NJ 1

ELANTH is the trade mark for Benlate — ELANCO PRODUCTS COMPANY
 DIVISION OF ELLI LILLY & COMPANY

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact, flush with water. Do not contaminate foodstuffs or feed.


FISH AND WILDLIFE CAUTION

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not spray where runoff is likely to occur. Do not contaminate anybody of water by direct application, cleaning of equipment, or disposal of wastes.

Do not reuse empty container. Destroy by burying with wastes or burning. Stay away from smoke or fumes.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

DESTROY BAG WHEN EMPTY. DO NOT REUSE.



PROFESSIONAL CRABGRASS AND POA ANNUA KILLER in late to early fall (mid-July to mid-September) prior to annual bluegrass (poa annua) germination. Chemicals that effectively control poa annua seeds may thin established poa annua and fine-leaved fescues. Reseeding should be delayed 12 to 16 weeks after application.

HOW TO APPLY:

Apply LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER evenly over the turf area without skips or misses. A more uniform application can be made by spreading half the required amount of this product over the area, and then applying the remaining half at right angles or crosswise. Both drop-type and rotary-type spreaders may be used to apply LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER.

Listed below are several suggested spreader settings. These should serve only as initial starting points for calibration since all spreaders will vary in their operation depending on the age, condition, particular model, and speed of operation.

APPROXIMATE SPREADER SETTINGS FOR LOFT'S PROFESSIONAL CRABGRASS AND POA ANNUA KILLER (for applying the contents of this bag in one pass over the turf)

SUGGESTED SPREADER SETTINGS

- CYCLONE
- O. M. SCOTT
- LAWN BEAUTY

