


WARFARIN CONCENTRATE

INTENDED FOR USE ONLY AS A RODENTICIDE. TO BE USED ONLY IN MANUFACTURING,
FORMULATING, MIXING OR REPACKING



ACTIVE INGREDIENTS:

Warfarin (3-(acetylbenzyl)-4-hydroxycoumarin) .5%

INERT INGREDIENTS

99.5%

TOTAL

100.0%

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

CAUTION: Keep away from humans, domestic animals, wildlife, and pets. If swallowed by humans, domestic animals, or pets, this material may reduce clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

SUGGESTED USE: Mix thoroughly one part of WARFARIN with 19 parts, by weight, of a suitable semi-permanent bait material such as corn meal, rolled oats, nut crumbs, dog food meal, poultry mash etc. Dogs and cats will actively search out the baits of this type. Therefore, baits must be placed so as to positively prevent feeding by these animals. A small amount, up to 5%, of a bland oil such as mineral or corn oil, or any other edible oil, may be used to prevent excessive dusting where the mixture has a tendency to dust.

For rat control place minimum 4 ozs. of mixed bait in locations frequented by rats, and maintain an uninterrupted supply for approximately 10 days. For control of mice, place tablespoon amounts of bait at 8-10 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days. Use no mouldy, decomposed or sour baits. Where a continuous source of infestation prevails, permanent bait stations should be used and baits replenished as needed. Baiting should continue until complete lack of feeding is noted. This should be from 5 to 14 days. Baits should be placed in locations which are generally inaccessible to humans, domestic animals, wildlife, and pets.

NET CONTENTS _____ Pounds

Distributed by:
LOBEL CHEMICAL CORP.
100 CHURCH STREET
NEW YORK, NEW YORK 10007