



CATTLE DUST

For the Control of Horn Flies, Lice, Fleas, Ticks and Feather Mites on Milk Cows, Livestock and Poultry.

ACTIVE INGREDIENTS:

- * Cyfluthrin 0.05%
- ** Triphenylethylene Pyrethroids 0.05%
- Pyrethrins 0.05%

INERT INGREDIENTS

- 1. Diatomaceous earth (2.2%) (pyrethrin-free)
- 2. Silica gel (2.2%) (pyrethrin-free)
- 3. Lignin (2.2%) (pyrethrin-free)
- 4. Cellulose (2.2%) (pyrethrin-free)
- 5. Magnesium silicate (2.2%) (pyrethrin-free)
- 6. Magnesium carbonate (2.2%) (pyrethrin-free)
- 7. Magnesium hydroxide (2.2%) (pyrethrin-free)
- 8. Magnesium stearate (2.2%) (pyrethrin-free)
- 9. Magnesium sulfate (2.2%) (pyrethrin-free)
- 10. Magnesium chloride (2.2%) (pyrethrin-free)
- 11. Magnesium bromide (2.2%) (pyrethrin-free)
- 12. Magnesium iodide (2.2%) (pyrethrin-free)
- 13. Magnesium nitrate (2.2%) (pyrethrin-free)
- 14. Magnesium phosphate (2.2%) (pyrethrin-free)
- 15. Magnesium sulfide (2.2%) (pyrethrin-free)
- 16. Magnesium selenide (2.2%) (pyrethrin-free)
- 17. Magnesium telluride (2.2%) (pyrethrin-free)
- 18. Magnesium borate (2.2%) (pyrethrin-free)
- 19. Magnesium fluoride (2.2%) (pyrethrin-free)
- 20. Magnesium oxide (2.2%) (pyrethrin-free)
- 21. Magnesium hydroxide (2.2%) (pyrethrin-free)
- 22. Magnesium carbonate (2.2%) (pyrethrin-free)
- 23. Magnesium silicate (2.2%) (pyrethrin-free)
- 24. Magnesium sulfate (2.2%) (pyrethrin-free)
- 25. Magnesium chloride (2.2%) (pyrethrin-free)
- 26. Magnesium bromide (2.2%) (pyrethrin-free)
- 27. Magnesium iodide (2.2%) (pyrethrin-free)
- 28. Magnesium nitrate (2.2%) (pyrethrin-free)
- 29. Magnesium phosphate (2.2%) (pyrethrin-free)
- 30. Magnesium sulfide (2.2%) (pyrethrin-free)
- 31. Magnesium selenide (2.2%) (pyrethrin-free)
- 32. Magnesium telluride (2.2%) (pyrethrin-free)
- 33. Magnesium borate (2.2%) (pyrethrin-free)
- 34. Magnesium fluoride (2.2%) (pyrethrin-free)
- 35. Magnesium oxide (2.2%) (pyrethrin-free)

NET WEIGHT:

CAUTION

KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.
DO NOT RE-USE CONTAINER
DESTROY WHEN EMPTY

HART - DELTA, INC.



Cattle Dust

FOR DAIRY CATTLE

For the control of Horn Flies on dairy cattle. Apply 2 ounces (two heaping tablespoonsful) on the neck, back and upper sides. Dust may be rubbed in lightly to insure that it has penetrated into the hair. Repeat application in three weeks if necessary. Do not apply within 5 hours of milking or during milking. See other cautions below. Do not treat calves under one month of age.

FOR BEEF CATTLE, HOGS AND OTHER LIVESTOCK

For the control of horn flies, lice, fleas and ticks. Apply 1 to 3 ounces per animal, sprinkling on the neck, back and upper sides. Dust may be rubbed in lightly to insure that it has penetrated into the hair. Repeat application in three weeks if necessary. Do not treat animals under 1 month of age.

FOR SHEEP

Apply dust to sheep after shearing in the same manner as for other animals. Will give temporary relief for ticks (keds) on sheep. Do not treat lambs one month of age or younger.

FOR POULTRY

To control feather mites on poultry, work well down into feathers, under neck and around the vent. Dust nest and litter to prevent reinfestation. Do not treat chicks under 8 weeks.

CAUTION

AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. Avoid contamination of feed and food-stuffs.

Do not apply within 5 hours of milking or during milking. Do not contaminate feed or water.

WARNING

This product will kill fish. Keep out of lakes, streams and ponds.