



12 500  
 EPA

**BEST AVAILABLE COPY**

**ACCEPTED**

SEP 1 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 6175-27

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS  
CAUTION

Herrn/If swallowed: Avoid contact with eyes or clothing.  
Avoid contact with food or feedstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms. Keep out of  
lakes, ponds, or streams. Do not contaminate water by  
cleaning of equipment or disposal of wastes.

STATEMENT OF PRACTICAL TREATMENTS

IF SWALLOWED: Drink a large amount of milk or water.  
DO NOT induce vomiting because of aspiration hazard.  
Call a Physician or Poison Control Center immediately.

IF INHALED: Remove victim to fresh air. Apply artificial  
respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash  
affected areas with soap and water.

IF IN EYES: Flush with plenty of water. See a Physician  
immediately.

STORAGE AND DISPOSAL

Store in a cool place out of reach of children. Do not  
reuse empty container. Wash empty container and put in  
trash collection.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

LOT NO.  
EPA REG NO 8175-27  
EPA EST NO 8175-LA-1

STABLE CONCENTRATE

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.  
For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

For use in the control of houseflies, hornflies, gnats,  
and mosquitoes in horse barns, stables, and tack rooms.

For use in the control of ticks on horses and ponies.

Manufactured By  
**HART-DELTA, INC.**  
5085 CHOCOMA DRIVE  
BENEFIT HOUSE, LA VERNE

Reconstitute with tap water. Prepare the sprayer and dilution of  
STABLE CONCENTRATE in the amount specified for correct use  
(about one quart per horse). Dilute STABLE CONCENTRATE  
should be used within 24 hours.

Reconstituted Solution

TO KILL AND REPEL HOUSEFLIES, HORNFIES, GNATS  
AND MOSQUITOES: Dilute one (1) part STABLE CONCENTRATE  
with nine (9) parts water or oil.

TO KILL AND REPEL HORSEFLIES, STABLE FLIES,  
DIPTERIDAE AND FACEFLIES: Dilute one (1) part STABLE  
CONCENTRATE with five (5) parts water or oil.

TO KILL ATTACHED TICKS: Dilute one (1) part STABLE CONCENTRATE  
with five (5) parts water or oil. Dilute directly on Tick  
with saturated cloth. Avoid contact with eyes or nostrils.

TO CONTROL FLIES, MOSQUITOES AND GNATS IN HORSE  
BARN, STABLE, AND TACK ROOMS: Dilute one (1) part  
STABLE CONCENTRATE with four (4) parts water or oil.

FOR APPLICATION ON HORSES AND PONIES: Brush animal  
thoroughly to remove excess dirt and dust. Using appropriate  
brush as directed during grooming (do not scrub) a full coat and  
rub over the hair. Apply by rubbing and against hair coat then  
rub over the hair. Give special attention to face, neck, legs,  
shoulders, and places where ticks are most often found. Avoid  
contact with eyes or nostrils. Apply to thoroughly moulted hair.  
Coat should not be interfered by light brushing following application.  
For best results apply STABLE CONCENTRATE daily.

FOR APPLICATION AS A SPACE SPRAY: Using appropriate  
dilution as directed above, apply as a fog or fine mist, directing  
spray toward the ceiling and upper corners, with the area to be  
treated being about 2 ounces per 1000 cubic feet of space. For  
best results, shut doors and windows before spraying and keep  
doors closed for ten to fifteen minutes. Vacate the area until  
vents to before reoccupying. Repeat application as necessary.