

MALATHION 50%
EMULSIFIABLE CONCENTRATE

Active Ingredients:	
Malathion*	50.0%
Methylated Naphthalenes	39.0%
Inert Ingredients:	
.....	11.0%
TOTAL	100.0%

* O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

CAUTION
Keep Out of the Reach of Children

Harmful if swallowed, inhaled, or absorbed through skin. Avoid prolonged breathing of spray mist. Avoid repeated or prolonged contact with skin. Wash with soap and water after using Malathion. Wash clothing with soap and water before re-use. Avoid contamination of feed and foodstuffs. Never re-use containers, destroy them when empty.

FIRST AID

EXTERNAL: If spilled on skin or clothing wash thoroughly with soap and water and change clothes. For eyes, flush with water for 15 minutes and get medical attention promptly.

INTERNAL: If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Repeat until vomit fluid is clear. CALL A PHYSICIAN IMMEDIATELY. SYMPTOMS of poisoning are pin-point pupils, drooling, lachrymation, labored breathing, vomiting, marked tremors, diarrhea, and convulsions. If these symptoms should occur after handling or using Malathion, call a PHYSICIAN IMMEDIATELY.

NOTE TO PHYSICIAN: If the above symptoms are present, atropine sulfate is the drug of choice. Seller makes no warranty of any kind, express or implied, concerning results obtained by the use of this material. Buyer assumes all risk of use or handling whether or not in accordance with any suggestions, recommendations or directions of seller.

DIRECTIONS FOR USE

The following are general recommendations for the use of Malathion 50% Emulsifiable Concentrate for home and garden use. THE RECOMMENDATIONS ARE BASED ON MAKING 1 GALLON OF SPRAY. SPRAY THOROUGHLY AND REPEAT APPLICATIONS AS NECESSARY.

ORNAMENTAL INSECTS

For use on flowers, shrubs and shade trees: To control aphids, birch leaf miners, flour-lined leaf bug, Japanese beetle adults, lace bugs, leafhoppers, mealybugs, tarnished plant bugs, thrips, Whitefly and European pine shoot moth; use 1 teaspoonful in 1 gallon of water.

To control bagworms, boxwood leaf miners, oak kermes, tent caterpillar, azalea scale, black scale crawlers, euonymus scale, fletcher scale, juniper scale, leaf scale, magnolia scale, oyster shell scale, scurfy scale, soft brown scale, and spider mites; use 2 teaspoonfuls in 1 gallon of water.

Use 4 teaspoonfuls in 1 gallon of water for the pine needle scale.

Injury may occur on certain ferns, and some species of crassula.

HOUSEHOLD INSECTS

Use 3 tablespoonfuls of this product in 1 quart of water, and apply as a coarse spray or brush on window sills, baseboards, drainboards, under sinks, and stoves, cracks and crevices, to control roaches, ants, spiders, silverfish, crickets, clover mites and earwigs.

Be careful not to contaminate food or utensils. Water base sprays may stain finished surfaces. Repeat application as necessary.

USDA Registration No. 6186-

DAMON CHEMICAL CO.

Box 480

Alliance, Ohio

Net Contents . . . 5 Gallons