

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 6190-2

DIRECTIONS:

Dilute 1 ounce of SANICO LF-34 to 3-5 gallons of water and sprinkle on 1000 square feet of grass.

SANICO LF-34 is easy to use. Best results are obtained by repeating this procedure every 10 days during the disease season. Even when applied at double the recommended strength, SANICO LF-34 does not injure the finest turf.

SANICO LF-34 goes into solution easily with water. Can be applied at all temperatures.

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

Antidote: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give two tablespoonfuls of Epsom Salt or Milk of Magnesia in water and plenty of milk or water. Have victim lie down and keep warm. Call a Physician Immediately. May Be Fatal If Swallowed! Do not get in eyes or on skin. Do not breathe spray mist. Wash thoroughly after handling.

This product is toxic to fish and wildlife. Keep out of all water supplies. Bury all wastes and containers in non-crop land away from water supply.

Danger: Wear a respirator when spraying, and wash contaminated clothing.

Do not take internally — if swallowed, induce vomiting immediately using mustard water, salt water, finger down throat or other methods. Keep patient warm to prevent shock.

EPA Reg. No. 6190-2

SANICO • 13143 Saticoy Street • North Hollywood, California 91605 •

SANICO

LF-34

a liquid fungicide

that is specially prepared for the prevention and control of brown patch, dollar spot, and pink patch.

DO NOT GRAZE TREATED AREAS.
DO NOT FEED CLIPPINGS TO LIVESTOCK.



**DANGER
POISON**



**KEEP OUT OF REACH OF CHILDREN —
See side panel for additional cautions.**

ACTIVE INGREDIENT:	
Cadmium Chloride _____	22%
INERT INGREDIENTS:	
Water _____	78%

**Net Contents
1 U.S. Gallon**

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 6190-2

DIRECTIONS:

Dilute 1 ounce of SANICO LF-34 to 3-5 gallons of water and sprinkle on 1000 square feet of grass.

SANICO LF-34 is easy to use. Best results are obtained by repeating this procedure every 10 days during the disease season. Even when applied at double the recommended strength, SANICO LF-34 does not injure the finest turf.

SANICO LF-34 goes into solution easily with water. Can be applied at all temperatures.

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

Sanico LF-34 is a liquid fungicide...
It is used for the prevention and control of brown patch, dollar spot, and pink patch on lawns and golf courses.
It is easy to use and does not injure the finest turf.
It goes into solution easily with water and can be applied at all temperatures.
It is registered under EPA Reg. No. 6190-2.

EPA Reg. No. 6190-2

SANICO • 13143 Saticoy Street • North Hollywood, California 91605 •

SANICO

LF-34

a liquid fungicide

that is specially prepared for the prevention and control of brown patch, dollar spot, and pink patch.

DO NOT GRAZE TREATED AREAS.
DO NOT FEED CLIPPINGS TO LIVESTOCK.



SANICO
POISON

KEEP OUT OF REACH OF CHILDREN —
See side panel for additional cautions.

ACTIVE INGREDIENT:	
Cadmium Chloride	22%
INERT INGREDIENTS:	
Water	78%

Net Contents
1 U.S. Gallon