

181

B-T-F

IODOPHOR SANITIZER*

601-12

"more concentrated"
for
sanitizing
GLASSES — DISHES — UTENSILS
in
Restaurants, Bars,
Hotels, Schools, Institutions

ACTIVE INGREDIENTS:

Butoxy polypropoxy polyethoxy ethanol-iodine complex (providing 1.6% titratable iodine)	12.54%
INERT INGREDIENTS	87.46%
	100.00%

*APPROVED UNDER THE REGULATIONS OF THE
WISCONSIN STATE BOARD OF HEALTH

EPA Reg. No. 6198-12-AA EPA, Est. No. 6198-MN-1
U.S. Patent No. 3,028,299

Net Contents
500 ml., 1/2 L. (16.9 fl. oz. or 1 Pint .9 oz.)

CAUTION: Keep out of reach of children.
(see additional cautions on back panel)

NATIONAL CHEMICALS Inc.,
Winona, MN 55987

ACCEPTED
6/98-12
JUL 23 1980
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the purpose
registered under
EPA Reg. No. 6198-12-AA

TD = 8.59 #/gal

473.2 ml 16 oz

354.9 ml 12 oz

8 oz

118.3 ml 4 oz

59.1 ml 2 oz

DIRECTIONS FOR MAKING SANITIZING SOLUTIONS (ODORLESS — TASTELESS)

Add B-T-F IODOPHOR to cool or lukewarm
water. Never use hot water.

1/4th oz. in 2 1/2 gal. water } 12.5 ppm
capful = 1/4 oz. } titratable iodine

1/5th oz. in 1 gal. water } 25 ppm
3/5th oz. in 3 gals. water } titratable iodine

Metric Equivalent: 1/10th oz. = 3 ml.
1/5th oz. = 6 ml. 1 gal. = 3.79 L.

NOTE: As long as solution is amber color,
there is sufficient iodine present
to sanitize. Make new solution
daily or when color fades.

RECOMMENDED PROCEDURE FOR WASHING & SANITIZING

1. Scrape & prewash glasses, dishes, etc.,
whenever possible.
2. Wash with a good detergent (T-D-C,
L-f-D, BLU MAGIK).
3. Rinse thoroughly with plain water.
4. Sanitize in B-T-F IODOPHOR solution
made to ppm concentration recom-
mended by Health Dept. Immerse
items for 2 minutes. Health require-
ments vary; so, check governing sani-
tary code for concentration and contact
time prescribed by local regulations.
5. Place sanitized items on rack or drain-
board to air dry. Never towel.

CAUTION: Avoid contact with food.
Harmful if swallowed. Avoid contact
with skin and eyes. In case of contact
with eyes, flush immediately with
water. Get medical attention.
RINSE EMPTY BOTTLE THOROUGHLY
WITH WATER AND DISCARD IT.

2013

BEST AVAILABLE COPY