

The following insertions are proposed on the attached label in the places where they are indicated by the appropriate single or double asterisk.

* "ALSO CAN BE USED ON GREENHOUSE LETTUCE, CUCUMBERS, TOMATOES, AND RADISHES. (See "LIMITATIONS FOR GREENHOUSE VEGETABLES" below)."

* "LIMITATIONS FOR GREENHOUSE VEGETABLES: 24 HOURS MUST BE PERMITTED TO ELAPSE BETWEEN TIME OF FINAL TREATMENT AND HARVESTING."

FLORA-FUME VAPONA[®]

DANGER

ANTIDOTES

If swallowed: If a child has ingested a small amount, repeat until vomit fluid is clear. Have a doctor examine child.

Call a physician immediately.

If on skin: Wash immediately with soap and water.

ATR-PNE-11-11-57

DANGER

Acute toxicity: Inhalation of vapors may cause irritation of the respiratory tract. Symptoms include coughing, sneezing, and difficulty breathing.

DO NOT breathe vapors.

Precautions: Do not use in enclosed spaces. Do not use near food or feedstuffs.

First aid: If inhaled, move to fresh air. If symptoms persist, seek medical attention.

Storage: Store in a cool, dry place. Keep away from heat and open flames.

Disposal: Dispose of according to local regulations. Do not pour down the drain.

USDA REGISTRATION NO. 6218-26

ACCEPTED

DANGER! KEEP OUT OF REACH OF CHILDREN!

POISON



SEE LEFT PANEL FOR ANTIDOTES AND PRECAUTIONS



VAPONA[®]

GREENHOUSE MISTING INSECTICIDE


Formulated Exclusively For Geiger Corporation, Harleysville, Pa

BALTIMORE



MARYLAND

ACCEPTED WITH COMMENTS

KEEP OUT OF REACH OF CHILDREN!
POISON
SEE INSTRUCTIONS FOR ANTIDOTES AND PRECAUTIONS 



E MISTING

FLORA-FUME VAPONA

DIRECTIONS FOR USE

PREPARATION OF GREENHOUSE

METHODS OF APPLICATION

Exclusively For Geiger Corporation, Harleysville, Pa

ORE



MARYLAND