

Summit

CAUTIONS

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc. where exposed food will be handled, with potable water. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

Physical Hazards

Do not store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.
Pesticide Disposal: Pesticide or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

- Container Disposal:** a) Containers of 30 gallons or more — reseal container and offer for reconditioning, or:
b) All containers — triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place.

General: Consult Federal, State or Local authorities for approved alternative procedures.

PYRETHRINS FOGGING CONCENTRATE II

For Use In Meat, Poultry, Egg,
and Rabbit Processing Plants

ACTIVE INGREDIENTS:

Pyrethrins	100%
Pyrethrins	2.0%
*Piperonyl butoxide, technical	2.5%
†N-octyl bicycloheptene dicarboximide	2.5%
Petroleum distillate	93.0%

*Equivalent to 2.0% (butylcarbityl) (6-propylpiperonyl) ether and 0.5% related compounds.
†MGK-264, Insecticide Synergist.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
See other cautions on side panel.

This synergized pyrethrins insecticide formulation is especially designed for use in Meat, Poultry, Egg and Rabbit Processing Plants. It may also be used in other indoor facilities where the insects shown on this label are present.

Insecticides for the Food Industry

BALTIMORE

A PRODUCT OF
Summit
CHEMICAL CO.

MARYLAND

ACCEPTED

24
MAY 5 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 612-45

PYRETHRINS FOGGING CONCENTRATE II

For Use In Meat, Poultry, Egg,
and Rabbit Processing Plants

ACTIVE INGREDIENTS:	100%
Pyrethrins	2.0%
*Piperonyl butoxide, technical	2.5%
†N-octyl bicycloheptene dicarboximide	2.5%
Petroleum distillate	93.0%

*Equivalent to 2.0% (butylcarbityl) (6-propylpiperonyl) ether and 0.5% related compounds.
†MGK-264, Insecticide Synergist.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
See other cautions on side panel.

This synergized pyrethrins insecticide formulation is especially designed for use in Meat, Poultry, Egg and Rabbit Processing Plants. It may also be used in other indoor facilities where the insects shown on this label are present.

BALTIMORE

A PRODUCT OF
Summit
CHEMICAL CO.

MARYLAND

ACCEPTED

MAY 5 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 6218-45

Summit

DIRECTIONS FOR USE (CONTINUED)

Because of the small micron size particles emitted from cold aerosol foggers, such as the LECO and MICRO-GEN, this dual synergized insecticidal formulation is extremely effective on those insects listed. The small particle size permits greater insect-insecticide contact than previously attainable by other methods of space spraying.

Prior to application, close all doors and windows. Turn off air-conditioning or ventilating systems. Cover or remove all food and food handling surfaces.

Space Spray to Kill Flies, Mosquitoes, Small Flying Moths and Gnats: Using a spraying device similar to the Micro-Gen or Mini Leco, apply 1/2 ounce per 1,000 cu. ft. of room space. The applicator should be centrally located or move around the room as necessary to achieve uniform distribution of the insecticide mist. Vacate premises and leave room closed for one hour. Ventilate room thoroughly prior to re-entry. Repeat as necessary for control.

Contact Spray to Kill Roaches, Silverfish, Waterbugs, Earwigs, Ants, Spiders, and Beetles in Sheltered Situations: With equipment described above, apply the spray into all cracks and crevices in woodwork, walls, floors, underneath, sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Do not allow spray to contact stored food. Repeat dosages as necessary for control. Leave the room closed for one hour. Do not remain in treated areas and ventilate area before re-entry.

NET CONTENTS

EPA REGISTRATION NO. 6218-45
EPA EST. 6218-MD-2

25