

USE DIRECTIONS:

DISINFECTION

For heavy duty cleaning and disinfecting and fungicidal activity against pathogenic fungi, use 4.5 ounces per gallon.

For best results, use FLAG GERMICIDE CLEANER with a cloth or mop and apply to walls, floors, table tops, and counters.

Sanitization

For light duty cleaning, sanitizing and deodorizing, use 2 ounces per gallon.

If used for sanitization of previously cleaned food contact items, limit the level to 200 ppm of quaternaries, (8 ounces per 5 gallons of water). A potable water rinse is required.

For extra heavy industrial ceaning and sanitizing, use a second application.

Do not mix with soap or anionic solutions.

FLAG GERMICIDE CLEANER is non-corrosive in use dilutions on inanimate hard surfaces.

PHENOL COEFFICIENTS:

Guaranteed	Minimum
S. aureus	14
S. typhosa	9



Suppliers of Top Quality Products to the San Joaquin Valley

GERMICIDE CLEANER
CLEANER DISINFECTANT
DEODORIZER FUNGICIDE

ACTIVE INGREDIENTS:

- n-Alkyl (60% C14, 30% C16, 5% C12 5% C18) dimethyl benzyl ammonium chlorides0.8%
- n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides0.8%
- Sodium Metasilicate2.4%
- Tetrasodium ethylenediamine tetraacetate1.0%
- Perfumes (Essential Oils)0.2%

INERT INGREDIENTS: 94.8%

DANGER 100.0%

KEEP OUT OF REACH OF CHILDREN.

See right side panel for additional precautionary and first aid statements.

EPA Reg. No. 6233-12

NET CONTENTS: _____

FLAG JANITORIAL SUPPLIES

2430 East McKinley

Fresno, CA 93703

MFG by Amador Chemical Corp.

Stockton, CA

REGISTERED

MAY 30 1975

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT

FOR ECONOMIC POISON REGISTERED UNDER NO. 6233-12

FLAG GEM cleaner ar purpose u sanitizing as directed remove m single, sir industrial shown on floors, tab motels, sc Not recd aeruginos For saniti processing limit the ounces pe water rins

Keep Out Causes se get in eyes and skin w if swallow

In case of skin with minutes. and wash reuse. If gelatin so drink larg physician

Probable r the use of circulatory convulsion Rinse emp and discar



Suppliers of Top Quality
Products to the San Joaquin Valley

GERMICIDE CLEANER

CLEANER DISINFECTANT
DEODORIZER FUNGICIDE

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides	0.8%
n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides	0.8%
Sodium Metasilicate	2.4%
Tetrasodium ethylenediamine tetraacetate	1.0%
Perfumes (Essential Oils)	0.2%
INERT INGREDIENTS:	94.8%

DANGER 100.0%

KEEP OUT OF REACH OF CHILDREN.

See right side panel for additional precautionary and first aid statements.

EPA Reg. No. 6233-12

NET CONTENTS: _____

FLAG JANITORIAL SUPPLIES

2430 East McKinley

Fresno, CA 93703

MFG by Amador Chemical Corp.
Stockton, CA

REGISTERED

MAY 30 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT

FOR PSEUDOMONAS AERUGINOSA REGISTERED
EPA UNDER NO. 6233-12

FLAG GERMICIDE CLEANER is a heavy duty cleaner and disinfectant designed for general purpose use where heavy duty cleaning and sanitizing is of prime importance. When used as directed, FLAG GERMICIDE CLEANER will remove most household dirt and grime in a single, simple application. When used for industrial cleaning, follow the use directions shown on the left side panel. For use on walls, floors, table tops, in industrial plants, offices, motels, schools, and homes.

Not recommended for Pseudomonas aeruginosa.

For sanitization of previously cleaned food processing equipment or food contact items, limit the active quaternaries to 200 ppm (8 ounces per 5 gallons of water). A potable water rinse is required.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.