


30
oz.
28
26
24
22
20
18
16
14
12
10
8
6
4



Net Contents 1 Quart

ACTIVE INGREDIENTS: n-alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides		42.0%
n-dialkyl (60% C14, 30% C16, 5% C12, 5% C18) methyl benzyl ammonium chlorides		8.0%
INERT INGREDIENTS		50.0%
		100.0%

No. 300040

E.P.A. Reg. No. 6284-5

DANGER

KEEP OUT OF REACH OF CHILDREN

See Cautions Elsewhere On Package

DISTRIBUTED BY

KIEFER **McNEIL**
DIVISION CORPORATION
999 Switzer Ave. A.V.A. Akron, Ohio 44311

AC
MAR - 17 - 1975
625

RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD.
THIS PRODUCT IS TOXIC TO FISH. DO NOT DISCHARGE EFFLUENT WHERE IT WILL DRAIN INTO LAKES, STREAMS OR PONDS.

TO CONTROL ALGAE IN SWIMMING POOLS

Provides economical protection against algae growth for at least 3 to 5 months in a 10,000 gallon swimming pool.

DIRECTIONS

1. Upon filling pool add 6 ounces of End-O-Alge for every 10,000 gallons of water.
2. Follow this by adding 1 1/3 ounces per 10,000 gallons every 7 days.
3. When algae growth is visible use 24 ozs. per 10,000 gallons of water. Clean pool when algae is dead. Drain pool, if necessary and refill. Repeat initial dose of 6 ozs. of End-O-Alge per 10,000 gallons of water. Container provides protection for over 52,000 gallons of water.
4. Use other chemicals as recommended on individual packages.

DANGER. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID. In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN. Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by the cautious intravenous injection of a short-acting barbituate drug.

KEEP OUT OF REACH OF CHILDREN

ACCEPTED WITH COMMENTS