

CPC CONCENTRATED-CHLORINE TABLETS

Slow dissolving, completely soluble CPC Concentrated-Chlorine Tablets take the work out of pool care by providing a continuous and economical system of chlorine disinfection. For effective bacteria and algae control, maintain an available chlorine level of 1.0 to 2.0 ppm as determined by appropriate test kit.

POOL START-UP (PRE-CONDITIONING) INSTRUCTIONS

When filling your new pool or refilling with fresh water at any time:

1. Super chlorinate pool water by adding CPC Pool Shock or CPC Granular Chlorine as prescribed on the labels.
2. Bring pool water pH to a range reading between 7.2 and 7.6 on your test kit and add CPC Water Conditioner to stabilize the pool water.
3. Start tablet feeding program.


TABLET FEEDING PROGRAM

FOR PROPERLY PRE-CONDITIONED POOLS

The number of tablets required for your pool may vary depending on temperature, wind, rain, pool usage and feeding method; float, submersible cage or skimmer.

6284-11
ACCEPTED

To start your tablet feeding program put 10 tablets in your feeder for every 10,000 gallons of pool water. TEST YOUR POOL FOR CHLORINE LEVEL DAILY and add tablets as required to maintain an available chlorine level of 1 to 2 ppm. DO NOT THROW TABLETS DIRECTLY INTO POOL. SPOTTING WILL OCCUR. FOR MAXIMUM PROTECTION SUPER CHLORINATE YOUR POOL WITH CFC POOL SHOCK EVERY TWO WEEKS AS PRESCRIBED ON THE LABEL.



DANGER

CORROSIVE

Causes eye damage. May cause burns on broken skin.

Irritating to nose and throat if inhaled. May be harmful or fatal if swallowed. Do not get in eyes, on skin or on clothing. Protect eyes when handling. Avoid breathing dust.

STRONG OXIDIZING MATERIAL. Contact with organic matter or any foreign material may result in fire or explosive decomposition. Moisture contamination or fire may liberate irritating and hazardous chlorine containing gases. Avoid contamination with combustible materials. Keep away from heat and moisture. Store in cool, dry, well-ventilated area. Keep container tightly closed when not in use. Use only dry, clean measures. Mix only with water. Rinse container thoroughly clean with water before discarding.

FIRST AID

Internal- feed cooked cereal or bread soaked in milk, followed by olive oil or cooking oil.

External- in case of contact, immediately flush eyes or skin with cold water for at least 15 minutes.

"CALL A PHYSICIAN IMMEDIATELY."

KEEP OUT OF REACH OF CHILDREN

Distributed by Kiefer-McNeil Akron, Ohio 44311

6284-11