



DANGER: KEEP OUT OF REACH OF CHILDREN See additional precautions and first aid on back panel

NO. 300-125

E.P.A. REG. NO. 6284-18

CPC POOL SHOCK FOR SHOCK TREATMENT OR SUPER CHLORINATION OF POOL WATER DIRECTIONS FOR SWIMMING POOL USE

Pool Shock is used for periodic shock treatments or super chlorination of water to eliminate a discomforting irritation due to formed chloramines in the pool water and to control algae.

Add 2 oz. of Pool Shock per 1,000 gallons of water by sprinkling it over the pool surface, or by sprinkling it into a bucket of water and pouring the dissolved solution into the pool. This procedure should be repeated every 2 weeks and should be done in the evening or at any time when the pool is free of bathers. Allow the available chlorine residual to drop to 2 ppm in properly stabilized water (1 ppm for unstabilized water) before entering the pool. Use Appropriate Test Kit to determine chlorine level.

Frequency of shock treatment depends on the extra demands for chlorine resulting from heavy bathing loads, accumulations of dust and dirt caused by storms, or continued high temperatures. For routine chlorine dosage follow the directions for using your regular chlorine product.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water.

DANGER

KEEP OUT OF REACH OF CHILDREN. MAY CAUSE CHEMICAL BURNS. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. MAY BE FATAL OR HARMFUL IF SWALLOWED. STRONG OXIDIZING AGENT. NEVER MIX WITH ANY MATERIALS OTHER THAN WATER. FIRE OR EXPLOSION MAY OCCUR IF THESE DIRECTIONS ARE NOT FOLLOWED. DO NOT REUSE CONTAINER. RINSE EMPTY CONTAINER THOROUGHLY WITH WATER BEFORE DISCARDING. USE ONLY A CLEAN DRY DISPENSER.

In case of fire, drench with large amount of water and cool surrounding areas with spray. Do not get in eyes or on skin. Store in the original container with CLOSED COVER in a COOL, DRY place. Dispose of spilled Pool Shock by immediately flushing with water.

FIRST AID

EXTERNAL — Flush skin or eyes with plenty of water. If skin irritation persists, get medical attention. For eyes, get prompt medical attention.

INTERNAL — Drink large quantities of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Distributed by KIEFER McNEIL

Akron, Ohio 44311