

JUL 20 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
OR ECONOMIC POISON REGISTER-
D UNDER NO. 6234-28

CPC GRANULAR CHLORINE II POOL START-UP (PRE-CONDITIONING) INSTRUCTIONS

When filling your new pool or refilling with fresh water at any time:

1. Super Chlorinate pool water by adding CPC Concentrated Granular Chlorine II at a rate of 10 oz. per 10,000 gallons of water. Alternate procedure: add CPC Pool Shock as prescribed on label. Allow the available chlorine residual to drop to 2 ppm (1 ppm for unstabilized water) before entering pool. Use appropriate test kit to check.
2. Bring pool water pH to a range reading between 7.2 and 7.6 on your Test Kit and add CPC Water Conditioner.
3. Start daily maintenance dosage.

DAILY MAINTENANCE DOSAGE FOR PROPERLY PRE-CONDITIONED POOLS

Test your pool water frequently during the day—apply the amount of CPC Granular Chlorine II required to maintain a test kit reading from 1 to 2 ppm available chlorine (approximately 1 to 2 ounces per 10,000 gallons per day). Maintain pH reading in a range from 7.2 to 7.6 on your test kit.

FOR MAXIMUM PROTECTION SUPER CHLORINATE (SHOCK) YOUR POOL WATER WITH CPC POOL SHOCK EVERY TWO WEEKS AS PRESCRIBED ON THE LABEL.

More frequent chlorine dosage may be required from time to time to meet extra demands for chlorine resulting from heavy bathing loads, accumulations of dust and dirt by storms, or continued high temperatures.

DANGER

Causes eye damage. May cause burns on broken skin. Dust irritating to nose and throat. May be harmful or fatal if swallowed. Do not get in eyes, on skin or on clothing. Protect eyes when handling. Avoid breathing dust.

STRONG OXIDIZING MATERIAL. Contact with organic matter or any foreign material may result in fire or explosive decomposition. Moisture contamination or fire may liberate irritating and hazardous chlorine containing gases. Avoid contamination with combustible materials. Keep away from heat and moisture. Store in cool, dry, well ventilated area. Keep container tightly closed when not in use. Use only dry, clean measures. Mix only with water. Rinse container thoroughly clean with water before discarding.

FIRST AID. Internal—feed cooked cereal or bread soaked in milk, followed by olive oil or cooking oil.

External—in case of contact, immediately flush eyes or skin with cold water for at least 15 minutes. "CALL A PHYSICIAN IMMEDIATELY."

DANGER: KEEP OUT OF REACH OF CHILDREN

Distributed by KIEFER McNEIL

Akron, Ohio 44311