

Another Laboratory

Controlled Product by



**DIRECTIONS FOR USE**

**COM-O-BRUSH** is a pre-mixed, ready-to-use spray which kills brush and woody vines. It is effective on a wide range of brush and woody vines. It is not intended for use on trees or shrubs.

**FOLIAGE TREATMENT** Foliar leaf stem sprays are used during the growing season on dense brush or woody vine growth such as Smilax, Greenbriers, and other woody vines. Apply **COM-O-BRUSH** after foliage is well developed using a drenching spray containing 3 to 4 quarts in 100 gallons of water. Coverage should be complete, wetting all parts of the plants, including leaves and stems to point of leaf fall. Applications are more effective when applied to actively growing plants. Higher volumes are necessary where the brush is very dense and over 8 to 10 feet high. Do not treat during periods of severe drought or in early fall when leaves have lost their healthy green color. Hard to kill species may need retreatment the following season.

**BASAL BARK TREATMENT** This treatment is not adapted for spraying dense stands of brush and woody vines. It can be used anytime of the year to obtain better kill of some species where dense brush and woody vines do not prevent proper application. Apply **COM-O-BRUSH** at the rate of 4 quarts in 25 gallons, 1 pint in 3 gallons of fuel oil, diesel oil, or kerosene, to the basal parts of stems, or tree trunks less than 6 inches in diameter, from ground line to a height of 12 to 16 inches, thoroughly wetting to run off all sides of the indicated area. Often a delayed response to treatment can be expected. Treated brush or trees preferably should not be cut for one year following treatment.

**STUMP TREATMENT** To prevent regrowth from cut stems and stumps, apply a solution of 4 quarts of **COM-O-BRUSH** in 25 gallons of water-free diesel oil, fuel oil, or kerosene, 1 pint in 3 gallons. Wet thoroughly the cut surfaces of the stump, concentrating the spray on the outside edges. Stumps of the stump and exposed roots until runoff to ground line is noticed. Best results are usually obtained when applied to freshly cut stumps with diameters in diameter or larger.

**FRUIT TREATMENT** For trees 4 inches in diameter or larger, use the foliage spray treatment. Cut a single half-circle of overlapping one-inch strips around the tree just above the fruit line. Treat the cut area with a solution of 2 quarts of **COM-O-BRUSH** in 25 gallons of fuel oil, diesel oil, or kerosene, 1 pint in 3 gallons.

ACCEPTED  
3-2-70  
6294-19

U.S.D.A. Registration No. 6294-19

# COM-O-BRUSH

## 5 LV

### BRUSH KILLER

(2,4-D/2,4,5-T Low-Volatile Emulsifiable Concentrate)

Contains 2.0 Pounds Each of 2,4-D and 2,4,5-T Acid Equivalent Per Gallon

**ACTIVE INGREDIENTS:**

Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid	34.0%
Isooctyl Ester of 2,4,5-Trichlorophenoxyacetic Acid	32.5%

**INERT INGREDIENTS:**

TOTAL	100.0%
-------	--------

\*Equivalent to 2,4-Dichlorophenoxyacetic Acid 22.5%

\*\*Equivalent to 2,4,5-Trichlorophenoxyacetic Acid 22.5%

**CAUTION** KEEP OUT OF REACH OF CHILDREN.  
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Quality Products of Distinction

COMET MANUFACTURING CORPORATION

1381 DALON DRIVE, N. E. • ATLANTA, GEORGIA

**GENERAL INFORMATION**

**COM-O-BRUSH BRUSH KILLER** is an emulsifiable concentrate which forms an emulsion in water and will readily mix with water-free diesel oil, fuel oil or kerosene.

**PREPARING THE SPRAY** Fill the spray tank with half the required amount of water or water-free diesel oil, fuel oil, or kerosene, add the recommended amount of **COM-O-BRUSH** emulsifiable concentrate with agitation, and continue filling the spray tank with water or petroleum oil in straight oil mixture; do not let water get into the **COM-O-BRUSH** itself or into the finished product.

**TREATING SMALL AREAS** One tablespoonful of **COM-O-BRUSH** in 1 1/2 gallons of water is about equal to 1 quart in 100 gallons of water.

**CAUTION** Avoid spray drift to susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. Spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less likely to drift. Although this ester is much less volatile than butyl or isopropyl esters, high temperatures may produce vapors after application which might injure some susceptible plants growing nearby.

Contamination of shallow fish-bearing waters at application rates recommended on this label may kill fish.

DO NOT graze treated areas by dairy animals within seven days after treatment.

DO NOT contaminate irrigation ditches or water used for domestic purposes.

DO NOT use the same spray equipment for applying other materials to plants, as injury will result.

DO NOT store near fertilizers, seeds, insecticides or fungicides.

DO NOT take internally. Avoid contact with skin, eyes or clothing. The following plants are included among those susceptible to **COM-O-BRUSH**:

- |             |              |
|-------------|--------------|
| Alder       | Gum          |
| Ash         | Honeysuckle  |
| Aspen       | Locust       |
| Birch       | Maple        |
| Box Elder   | Oak          |
| Brambles    | Osage Orange |
| Buckbrush   | Salmonberry  |
| Wild Cherry | Poison Ivy   |
| Elderberry  | Palmetto     |
| Elm         | Sumac        |
| Greenbrier  | Willow       |
| Wild Grape  |              |

**WARRANTY AND LIMITATION OF DAMAGES**

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

ACCEPTED WITH RESERVE