

DIRECTIONS FOR USE

TIME OF APPLICATION Best results are obtained when CMC WEED KILLER is used on weeds that are young and in a rapidly growing condition. Applications of lower rates to susceptible annual weeds will be satisfactory, but for perennial weeds and other conditions where weed kill is difficult, use higher rates. When used as a selective spray in crops, the stage of growth of the crop must be considered. Some hardy plants and weeds are hard to kill and repeat applications may be necessary.

SMALL GRAIN CROPS Wheat, Oats, Barley, Rye. Apply CMC WEED KILLER at the rate of 5 to 6 pints per acre on small grains when fully tillered or stalked, 4 to 8 inches tall, but before head emerges from the boot. Injury will be likely to occur if this product is applied earlier than tiller or later than boot. DO NOT use on grains undersown with legumes such as alfalfa or clovers, except where some legume injury can be tolerated. DO NOT apply when grain is headed.

DO NOT forage or graze treated grain fields within two weeks after treatment with 2,4-D.

AERIAL APPLICATION To small grains, Wheat, Oats, Barley, and Rye and Rice. Use CMC WEED KILLER at the rate of 5 to 6 pints in sufficient water to insure adequate coverage.

RICE: In the South and Southwest apply from 1 to 1 1/2 gal per acre in 5 to 10 gallons of water 7 to 10 weeks after planting, or when the rice is fully tillered but not yet in the boot. In California, apply 1 to 1 1/2 gal per acre 55 to 65 days after planting, when the rice is through the water and well established, but not after seed stalks emerge. For difficult weeds and where daily temperatures do not exceed 95 F., 2 gals per acre may be used.

CORN: PREEMERGENCE Use CMC WEED KILLER at the rate of 1 1/4 to 2 1/2 gal in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting, but before corn emerges. DO NOT apply to light sandy soils.

POSTEMERGENCE apply at the rate of 5 pints per acre. When spraying corn above 10 inches in height, use nozzle extensions "corn drops", directing the spray down the base of the corn plant. DO NOT apply from tasseling to dough stage.

SORGHUM: PREEMERGENCE Use spray at the rate of 5 pints per acre on plants 6 to 12 inches high when secondary roots are well established. DO NOT apply from flowering to dough stage. Use drop nozzle when crop is over 10 inches tall.

SUGAR CANE Apply at the rate of 1 1/4 to 2 1/2 gal per acre as a spring preemergence treatment or through layby.

PASTURES To control many broadleaved weeds in pastures, meadows, and rangelands, use 1 1/2 to 2 1/2 gal per acre of CMC WEED KILLER in sufficient water to provide for uniform application. Treat pastures when weeds are growing actively. DO NOT apply to recently seeded pastures until grass is well established. DO NOT apply when grass is in boot or milk stage. Most legumes are usually injured or killed at the rates recommended. DO NOT graze treated areas by dairy animals within seven days after treatment.

LAWNS AND GOLF FAIRWAYS Apply 2 to 3 pints of CMC WEED KILLER per acre in sufficient water to obtain uniform coverage. DO NOT apply to St. Augustine grass, bermuda, bent grasses or to freshly seeded turf. Grasses have become well established. Most weeds are usually injured or killed.

FOR TREATING SMALL AREAS One tablespoon of this product in 1 1/2 gallons of water is about equal to one quart in 100 gallons of water.

GENERAL WEED CONTROL On vacant lots, roadsides and along fence rows use 2 1/2 to 4 gals of CMC WEED KILLER in 100 gallons of water per acre. Thoroughly wet all foliage to run off.

NOTE When stored at temperatures below freezing it may be necessary to warm up to at least 45 F. and shake thoroughly before using.



E.P.A. Registration No. 6294-23



WEED KILLER

2,4-D

Contains 0.8 Pounds 2,4-D Acid Equivalent Per Gallon

ACTIVE INGREDIENTS:	
Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic Acid*	10.0%
INERT INGREDIENTS:	
	90.0%
TOTAL	100%
*Equivalent to 2,4-Dichlorophenoxyacetic Acid 8.3%	

**KEEP OUT OF REACH OF CHILDREN.
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS
APPLY THIS PRODUCT ONLY AS
SPECIFIED ON THIS LABEL
HARMFUL IF SWALLOWED**

CAUTION

Quality Products of Distinction

NET VOL. GAL.

COMET MANUFACTURING CORPORATION

1381 DALON DRIVE, N.E. • ATLANTA, GEORGIA

GENERAL INFORMATION

CMC WEED KILLER is recommended for controlling Arr. 4 Broad, Dandelion, Docks, Herb, Goldenrod, Lambsquarters, Mustards, Peppergrass, Pigweed, Plantains, Ragweed, Shepherds-purse, Thistles, Wild Carrot, Wild Morning-glory, Wild Radish, and many other 2,4-D-susceptible broadleaved weeds in crops. The product also can be used on lawns, golf courses, parks, along highways, vacant lots and other areas without injury to most grasses.

PREPARATION OF THE SPRAY Fill the spray tank with half the required amount of water, then add the recommended amount of CMC WEED KILLER with agitation and continue filling the spray tank with water. Use enough water per acre to insure uniform coverage. The amount of water required for lawns, home applications may vary from 5 to 25 gallons per acre. For high volume application 100 gallons or more of water will be needed for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

CAUTION

- AVOID contact with skin, eyes or clothing.
- DO NOT store near fertilizers, seeds, insecticides or fungicides.
- DO NOT use the same spray equipment for applying other material to plants, or injury may result.
- DO NOT contaminate irrigation ditches or water used for domestic purposes.
- DO NOT contaminate water by cleaning of equipment or disposal of wastes.
- DO NOT reuse empty drum. Return to drum manufacturer or destroy by perforating or crushing and burying in a safe place away from water supplies.
- Exercise caution with regard to St. Augustine Grass and Dichroa, as possible injury might occur.
- Although CMC WEED KILLER is non-volatile, care must be used to prevent spray drift from coming into direct contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton and other desirable plants sensitive to 2,4-D. Coarse sprays are less likely to drift.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER be liable for any breach of warranty except the purchase price of the material as to which a claim is made.

ACCEPTED
Aug 19, 1974
UNITED STATES FEDERAL INSECTICIDE AND FUNGICIDE AGENCY
OFFICE OF THE REGISTERED CHEMIST
6294-23