

11/11/77

DIRECTIONS FOR USE

TIME OF APPLICATION: Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.**

WEEDS: Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.**

DO NOT apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.**

AERIAL APPLICATION: Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.**

RICE: In the South and Southwest apply from 1 1/2 to 2 1/2 pints per acre in 5 to 10 gallons of water 7 to 10 weeks after planting, or when the rice is fully tilled but not yet in the boot. In California, apply 1 1/2 to 2 pints per acre 55 to 65 days after planting when the rice is through the water and well established, but not after seed stalks emerge. For control of weeds and other daily for pastures, this product is 95% effective when used.

CORN: PRE-EMERGENCE: Use SUPERCID WEED KILLER at the rate of 1 to 2 quarts in soft cast water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting, but before corn emerges. **DO NOT** apply to light sandy soils.

POST-EMERGENCE: Apply at the rate of 1 pint per acre. When spray no more than 10 inches high. Use a nozzle extending 18 inches above the crop. Apply directly to the base of the plant. **DO NOT** apply to corn with seedling stage.

SORGHUM: POST-EMERGENCE: Apply at the rate of 1 pint per acre on plants 4 to 12 inches high. Apply when weeds are well established. **DO NOT** apply to corn. For control of weeds and other daily for pastures, this product is 95% effective when used.

SUGAR CANE: Apply at the rate of 2 to 4 quarts per acre.

PASTURES: Apply at the rate of 2 to 3 pints per acre of SUPERCID WEED KILLER in soft cast water to provide for uniform coverage. For control of weeds and other daily for pastures, this product is 95% effective when used. **DO NOT** apply to pastures with seedling stage. For control of weeds and other daily for pastures, this product is 95% effective when used.

ORNAMENTALS: Apply at the rate of 1 to 3 quarts per acre. For control of weeds and other daily for pastures, this product is 95% effective when used.

PRECAUTIONS: Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.**

GENERAL INFORMATION: Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.**

NOTES: Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.** Apply SUPERCID WEED KILLER to weeds in the following crops: **CORN, SOYBEANS, RICE, SUGAR CANE, PASTURES, AND ORNAMENTALS.**

Another Laboratory



SUPERCIDE CC WEED

Contains 4.0 Pounds 2,4-D /

ACTIVE INGREDIENTS:
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid

INERT INGREDIENTS:

TOTAL

*Equivalent to 2,4-Dichlorophenoxyacetic Acid

CAUTION KEEP OUT C
SEE SIDE PA
HARMFUL IF


NET CONTENTS

Quality Products

COMET MANUFACTURING

1381 DALON DRIVE, N. E.

Controlled Product by



U.S.D.A. Registration No. 6294-24

CONCENTRATED KILLER

acid Equivalent Per Gallon

oxyacetic Acid*	49%
.....	51%
.....	100%

acid 41%


REACH OF CHILDREN.
NEED FOR ADDITIONAL CAUTIONS
SWALLOWED

..... GALS.

of Distinction

NG CORPORATION

• ATLANTA, GEORGIA



GENERAL INFORMATION

SUPERCIDE WEED KILLER is recommended for controlling Arrow-Head, Dandelion, Dock, Henbit, Goldensrod, Lambsquarters, Mustards, Peppergrass, Pigweed, Plantains, Ragweed, Shepherds-purse, Thistles, Wild Carrot, Wild Morning-glory, Wild Radish, and many other 2,4-D susceptible broadleaved weeds in crops. This product also can be used on lawns, golf courses, parks, along highways, vacant lots and other areas without injury to most grasses.

PREPARATION OF THE SPRAY Fill the spray tank with half the required amount of water, then add the recommended amount of SUPERCIDE WEED KILLER with agitation and continue filling the spray tank with water. Use enough water per acre to give uniform coverage. The amount of water required for low volume applications may vary from 5 to 25 gallons per acre. For high volume application, 100 gallons or more of water will be needed for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

CAUTION

Avoid contact with skin, eyes or clothing.

DO NOT store near fertilizers, seeds, insecticides, or fungicides.

DO NOT use the same spray equipment for applying other materials to plants, as injury may result.

DO NOT contaminate irrigation ditches or water used for domestic purposes.

Exercise caution with regard to St. Augustine Grass and Dichondra, as possible injury might occur.

Although SUPERCIDE WEED KILLER is non-volatile, care must be used to prevent spray drift from coming into direct contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants sensitive to 2,4-D. Coarse sprays are less likely to drift.

Do not reuse empty drums. Return to drum rental firm or dealer by perforating or crushing and burying in a safe place away from water supplies.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.