

CAUTION: Do not inhale. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

GENERAL INFORMATION: TIGER Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its phytotoxic properties are quickly inactivated on contact with soil. It is a combination of herbicide and surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young weeds at air temperatures above 70 degrees F.

MIXING INSTRUCTIONS: TIGER Herbicide must be thoroughly dissolved. Fill the spray equipment reservoir about half full with water and add the required amount of herbicide with agitation. Finish filling the reservoir with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible and discard in a safe place.

DIRECTIONS FOR USE:

COTTON: TIGER Herbicide is useful as a directed application in cotton for post-emergent control of weeds and grasses, such as Johnsongrass, nutsedge, Dallisgrass, cocklebur, barnyard-grass, ragweed, sandbur and puncture vine. It should be applied at the rate of 2 quarts per acre.

Mix 5 quarts of TIGER Herbicide in 100 gallons of water. Do not add any surfactant to the spray solution. Apply spray solution at 40 gallons per acre.

HILL AP LENTY

TIGER BRAND


A HERBICIDE FOR SELECTIVE POST EMERGENT WEED CONTROL

Active Ingredient: Monosodium Acid Methanearsonate* 34.8%
Inert Ingredients: 65.2%
Total Arsenic (as elemental) all in water soluble form 16.1%
*Product contains 4 lbs. MSMA per gallon

CAUTION KEEP OUT OF REACH OF CHILDREN
Net Contents _____ Gallons

NON-WARRANTY: Adkins-Phelps Company makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Manufactured For
ADKINS-PHELPS COMPANY
CHEMICAL DIVISION
NORTH LITTLE ROCK, ARKANSAS



Directions For Use - Cont'd.
Weed control is most effective at temperatures above 70 degrees F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre. Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage. A second application may be required during humid growing conditions to control perennials, such as Johnsongrass and nutsedge. Apply the second application 1 to 3 weeks later. Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally. APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO FIRST BLOOM. DO NOT APPLY AFTER FIRST BLOOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK OR GRAZE TREATED AREAS.

NON-CROP: TIGER Herbicide is useful for control of Johnsongrass, nutsedge, barnyardgrass and other weeds on drainage ditchbanks, rights-of-way, fence rows, storage yards and similar non-crop areas. Mix TIGER Herbicide at the rate of 2 to 5 quarts in 100 gallons of water. Spray unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs re-apply as required. DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR BY ANIMALS, WILDLIFE AND AQUATIC LIFE, OR FOR IRRIGATION PURPOSES.

LAWN AND ORNAMENTAL TURF: TIGER Herbicide is useful for control of Dallisgrass, sandbur, bahiagrass, nutsedge, barnyardgrass, chickweed and wood sorrel with little or no injury to good lawn grasses. On new lawns do not treat until after three mowings. Good grasses may be temporarily discolored. Zoysia, bluegrass and bermuda are quite tolerant. Do not use on St. Augustine or centipede. Mow lawns 1 to 1 1/2 inches high before treatment. Mix 2 fluid ounces (4 tablespoons) of TIGER Herbicide in 5 gallons of water and apply to an area of 1,000 sq. ft. Spray thoroughly to wet all undesirable plants.

Repeat applications 10 to 14 days apart, may be needed for good control.

TIGER Herbicide is manufactured for
Adkins-Phelps Company
North Little Rock, Arkansas