

Phytar 535

Herbicide

July 6, 1973
6308-50


Sodium Cacodylate
Liquid Plus
Surfactant

For General
Post-Emergent
Weed Control

ACTIVE INGREDIENTS:
Sodium Cacodylate 28.05%
Dimethylarsinic Acid
(Cacodylic Acid) 4.81%
INERT INGREDIENTS: 67.14%
Total Arsenic (as elemental)
all in water soluble
form 15.80%
Product contains 25 lbs.
cacodylic acid equivalent
per gallon.

CAUTION:
Keep out of the
reach of children

Read entire label
before using
this product



SPECIMEN LABEL

CAUTION: Keep Out of the Reach of Children.

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

WARRANTY - CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Ansul makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Ansul or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.

GENERAL INFORMATION: PHYTAR 535 Herbicide is useful for general post-emergent weed control. It is non-selective in its action. Its phytotoxic properties are quickly inactivated on contact with soil. It is a combination of herbicide and surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds. It produces top-kill only so repeat applications are required for seasonal weed control of perennials.

MIXING INSTRUCTIONS: PHYTAR 535 Herbicide is completely water soluble. Fill the spray equipment tank about half full with water and add the required amount of herbicide with agitation. Finish filling the tank with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period. Although PHYTAR 535 Herbicide is only moderately corrosive, do not use in galvanized or aluminum equipment.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and a detergent, crush if possible, and discard in a safe place.

DIRECTIONS FOR USE:

NON-CROP: PHYTAR 535 Herbicide is useful for general weed control on drainage ditches, rights-of-way, along sidewalks, driveways and fences, around buildings and ornamentals, and similar non-crop areas. Mix 6 to 12 pints of PHYTAR 535 Herbicide in 100 gallons of water. Spray unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

Do not apply directly, or allow spray to come in contact with foliage, green bark, grafted unions, scuffed, damaged or broken bark when spraying around trees and ornamentals.

Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life or for irrigation purposes. Do not graze treated areas to livestock.

LAWN RENOVATION: Mow lawn to be renovated about 1 inch high before treatment. Mix fluid ounces of PHYTAR 535 Herbicide in 4 gallons of water and apply to 1,000 sq. ft. Cover foliage thoroughly, preferably on a warm sunny day. Foliage will usually turn brown in about five days. Bermudagrass and other deep-rooted perennials, or uncovered foliage, may require second treatment. Reapply five days later if green areas remain.

When foliage is all brown, matted areas should be raked and bare areas scarified. About five days after last treatment, water thoroughly and then seed at rate recommended for new lawns. Seed must be raked into soil. Keep seeded areas well watered.

Keep children and pets off treated areas until after first rain or sprinkling following treatment. Do not track from treated to untreated areas.

PHYTAR 535 Herbicide is manufactured by The Ansul Company, Marinette, Wisconsin 54142.

PHYTAR, ANSUL are registered trademarks of The Ansul Company.

Net Contents 5 Gallons
Form No.

EPA Reg. No. 6308-50



THE ANSUL COMPANY
MARINETTE, WISCONSIN