

Anorack®

Herbicide

Controls Seedling Grasses
and Weeds in Soybeans
and Peanuts


ACTIVE INGREDIENTS:

Sodium N-1-Naphthylphthalamate* 22.3%
Dinoseb, sodium salt (sodium 2-sec-butyl-4,6-dinitrophenate)** 11.5%

INERT INGREDIENTS: 66.2%

*Equivalent to 20.8% N-1 naphthylphthalamic acid.
**Equivalent to 10.5% 4,6 Dinitro o sec-butylphenol.
Each gallon contains two pounds of active NPA and one pound of active DNBP.

DANGER:
Keep out
of the reach
of children



DANGER: May be fatal if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Do not wear contaminated clothing. Do not breathe spray mist. Keep away from heat and open flame. Not for use or storage in or around the home.

This product is toxic to fish. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

SYMPTOMS OF POISONING: Excessive fatigue, sweating, thirst and fever. If symptoms of poisoning develop from any type of exposure, SEND FOR A PHYSICIAN.

FIRST AID: Have patient lie quiet in coolest spot available. If feverish, cool with cold compresses or by immersion in cool water.

ANTIDOTE: If swallowed, SEND FOR A PHYSICIAN. Induce vomiting by giving an emetic such as 2 tablespoons of table salt in a glass of warm water; repeat until vomit fluid is clear, then give 2 teaspoons of baking soda in a glass of water. Treat as in FIRST AID, above. **IF SPLASHED IN EYES,** immediately flush eyes with plenty of water for at least 15 minutes and get medical attention.

IF SPILLED ON SKIN, immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If symptoms of poisoning develop, send for a physician and treat as in FIRST AID, above. Discard contaminated clothing and shoes, or clean them thoroughly before re-use. Staining may result if contaminated articles are allowed to contact carpets, upholstery, etc.

NOTE TO ATTENDING PHYSICIAN: Active ingredient is a metabolic stimulant. Treat symptomatically.

CONTAINER DISPOSAL: Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Keep out of smoke.

WARRANTY - CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansol Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences

ALL SUCH RISKS ARE ASSUMED

BROADLEAF WEEDS: F. weed, lambsquarters, ragweed, leaf, cocklebur, bindweed, mustards, shephard's purse, carpetweed, morningglory, (lice) and teasweed (prickly

PREPARATION OF SOIL: treated should be well prepared from stubble and been plowed the soil closed up by harrow or roller or slightly raised bed. If replanting necessary after applied, thoroughly disk replanting.

METHOD OF APPLICATION: uniform coverage of the for good weed control. A spray at planting time using mounted sprayer or a separate planting operation. The applied up to the time of "cracking-time" on peanut plants emerge on soybean

DIRECTIONS FOR USE: recommended amount of tank, add the required amount then add the remainder tank while mixing. Stir until completely mixed. sprayer be calibrated according to crop.

PEANUTS: Use ANCR overall spray (broadcast) just as the peanuts begin or "cracking-time". This 7 days after planting. H be made from planting majority of the peanut "cracking-stage".

RECOMMENDED DOSAGE: **PEANUTS:** Use ANCR quarts in 20 to 40 gallons an overall spray (broadcast) may be used over bands as desired. The disturbed after planting with treatment is used


Amount of ANCRACK Use for Band Treatment

Width of Band	24 inches
12"	6 pints
14"	7 pints

Width of Band	32 inches
12"	4 1/2 pints
14"	5 1/2 pints


••Equivalent to 10.5% 4,6-Dinitro-o-sec-butylphenol.
Each gallon contains two pounds of active NPA
and one pound of active DNBP.

DANGER:
Keep out
of the reach
of children



POISON

See side panel
for antidote
and other
warning
statements



REGISTERED TRADE MARK

and wash skin with soap and water.
If symptoms of poisoning develop, send for a
physician and treat as in FIRST AID, above.
Discard contaminated clothing and shoes, or
clean them thoroughly before re-use. Staining
may result if contaminated articles are allowed
to contact carpets, upholstery, etc.

NOTE TO ATTENDING PHYSICIAN: Active
ingredient is a metabolic stimulant. Treat
symptomatically.

CONTAINER DISPOSAL: Rinse equipment and
containers and dispose of wastes by burying in
non-crop lands away from water supplies. Con-
tainers should be disposed of by punching holes
in them and burying with wastes, or by burn-
ing. Keep out of smoke.

WARRANTY - CONDITION OF SALE:
DIRECTIONS FOR USE of this product are
based on field use and tests believed reliable
and should be followed carefully. It is however
impossible to eliminate all risks associated with
use of this product. Because such factors as
weather conditions, foreign material and man-
ner of use for application are all beyond the
control of The Ansul Company or the Seller of
this product, such things as crop injury, inef-
fectiveness or other unintended consequences
may result. **ALL SUCH RISKS ARE ASSUMED
BY THE BUYER.**

Ansul warrants that this product conforms to
the chemical description on the label and is
reasonably fit for the purposes referred to in
the directions for use as modified by the above.
Ansul makes no other warranties, express or
implied, including **FITNESS or MERCHANT-
ABILITY.** In no case shall Ansul or the Seller
be liable for consequential, special or indirect
damages resulting from the use or handling of
this product. The foregoing is a condition of
sale by The Ansul Company and is accepted
as such by the Buyer.

GENERAL DIRECTIONS: Liquid ANCRACK
is a selective herbicide combination for the con-
trol of the following annual grasses and broad-
leaf weeds:

ANNUAL GRASSES: Foxtail, crabgrass, barn-
yardgrass, goosegrass, stinkgrass, sandbur, water-
grass, spangletop, windmillgrass, cupgrass and
johnsongrass (from seed).



THE ANSUL COMPANY, MARINETTE WISCONSIN

overall spray (broadcast) or
just as the peanuts begin to
or "cracking-time". This is
7 days after planting. How-
ever, it may be made from planting time
majority of the peanut plant
"cracking-stage".

RECOMMENDED DOSAGE
PEANUTS: Use ANCRACK
quarts in 20 to 40 gallons of
an overall spray (broadcast)
ment may be used over the
bands as desired. The soil sh-
turbled after planting when
treatment is used.

Amount of ANCRACK (per
Use for Band Treatment of


Width of Band	24 inches	Row of Band
12"	6 pints	5
14"	7 pints	6

Width of Band	32 inches	Row of Band
12"	4½ pints	4
14"	5½ pints	5

Use in 12-20 gallons of water.
Note 1. Plant on a flat or
Note 2. Space seed in row
(Spanish) or 4 inches
and plant at least
Note 3. May be used in the
following the use
incorporated herbicide.

Do not allow spray to contact
Crop Plants not recommended
injury may result.

CONTAINER STORAGE
ANCRACK is normally a solid
crystallize in the cold (32-40°F.)
warming (75-125°F.) will re-
Mild agitation, such as rolling
stirring before use is recom-
Avoid contamination of food
fertilizers, seeds, plants and
a cool dry location.



DANGER: May be fatal if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Do not wear contaminated clothing. Do not breathe spray mist. Keep away from heat and open flame. Not for use or storage in or around the home.

This product is toxic to fish. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

SYMPTOMS OF POISONING: Excessive fatigue, sweating, thirst and fever. If symptoms of poisoning develop from any type of exposure, SEND FOR A PHYSICIAN.

FIRST AID: Have patient lie quiet in coolest spot available. If feverish, cool with cold compresses or by immersion in cool water.

ANTIDOTE: If swallowed, SEND FOR A PHYSICIAN. Induce vomiting by giving an emetic such as 2 tablespoons of table salt in a glass of warm water; repeat until vomit fluid is clear, then give 2 teaspoons of baking soda in a glass of water. Treat as in FIRST AID, above. **IF SPLASHED IN EYES,** immediately flush eyes with plenty of water for at least 15 minutes and get medical attention.

IF SPILLED ON SKIN, immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If symptoms of poisoning develop, send for a physician and treat as in FIRST AID, above. Discard contaminated clothing and shoes, or clean them thoroughly before re-use. Staining may result if contaminated articles are allowed to contact carpets, upholstery, etc.

NOTE TO ATTENDING PHYSICIAN: Active ingredient is a metabolic stimulant. Treat symptomatically.

CONTAINER DISPOSAL: Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Keep out of smoke.

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, inef-

BROADLEAF WEEDS: Florida pusley, chickweed, lambsquarters, ragweed, purslane, velvet-leaf, cocklebur, bindweed, ground cherry, mustards, shephard's purse, pigweed, quickweed, carpetweed, morningglory, beggarweed (beggarlice) and teasweed (prickly sida).

PREPARATION OF SOIL: The area to be treated should be well prepared and as free as possible from stubble and litter. If the land has been plowed the soil clods should be broken up by harrow or roller wheel. Plant on a flat or slightly raised bed. If weather factors make replanting necessary after ANCRACK has been applied, thoroughly disk the area prior to replanting.

METHOD OF APPLICATION: Thorough and uniform coverage of the soil surface is essential for good weed control. Apply ANCRACK as a spray at planting time using either a planter-mounted sprayer or a separate sprayer after the planting operation. The chemical may be applied up to the time of crop emergence or "cracking-time" on peanuts but before the crop plants emerge on soybeans.

DIRECTIONS FOR USE: Put one-half the recommended amount of water in the spray tank, add the required amount of ANCRACK, then add the remainder of the water to the tank while mixing. Stir or use by-pass agitation until completely mixed. It is essential that the sprayer be calibrated accurately prior to use on crop.

PEANUTS: Use ANCRACK Herbicide as an overall spray (broadcast) or band application, just as the peanuts begin to crack the ground, or "cracking-time". This is normally from 5 to 7 days after planting. However, application can be made from planting time up to the time the majority of the peanut plants are in the "cracking-stage".

RECOMMENDED DOSAGE RATE FOR

PEANUTS: Use ANCRACK at the rate of 6 quarts in 20 to 40 gallons of water per acre as an overall spray (broadcast application). Treatment may be used over the entire area or in bands as desired. The soil should not be disturbed after planting when the "cracking-time" treatment is used.

Amount of ANCRACK Herbicide Per Acre to Use for Band Treatment of Peanuts

Width of Band	Row Spacing		
	24 inches	28 inches	30 inches
12"	6 pints	5 pints	4 1/2 pints
14"	7 pints	6 pints	5 1/2 pints

Width of Band	Row Spacing		
	32 inches	34 inches	36 inches
12"	4 1/2 pints	4 1/2 pints	4 pints

EQUIPMENT CLEANING: Thoroughly rinse spray tank, hose and nozzles with water after application. Use ANCRACK in metal tanks. It is difficult to remove from wooden tanks. ANCRACK can be removed from metal tanks by washing thoroughly with ammonia water and rinsing several times with water containing a good detergent.

DO NOT USE "ANCRACK" HERBICIDE ON SILT LOAM SOILS OF EXTREMELY FINE TEXTURE.

SOYBEANS: Use ANCRACK as an overall spray (broadcast), or band application, just after the soybeans are planted but before the soybean plants emerge from the soil.

RECOMMENDED DOSAGE RATE FOR SOYBEANS: Use ANCRACK at the rate of 6 quarts in 30 to 40 gallons of water per acre as an overall spray (broadcast application). Treatment may be applied over the entire area or in bands as desired.

Amount of ANCRACK Herbicide Per Acre to Use for Band Treatment of Soybeans

Width of Band	Row Spacing	
	28 inches	32 inches
12"	5 pints	4 1/2 pints
14"	6 pints	5 1/2 pints

Width of Band	Row Spacing	
	36 inches	40 inches
12"	4 pints	3 1/2 pints
14"	4 1/2 pints	4 1/4 pints


Use in 9 gallons of water per 40 inch row, to 13 gallons of water per 28 inch row.

Note 1. In the black soil areas of western Indiana, southern Minnesota, northern Iowa and Illinois, the dosage of ANCRACK suggested in the chart above should be increased by approximately one-third.

Note 2. Soybean seed should be well covered with soil. Do not use on shallow planted or exposed seed.

Note 3. If heavy rains immediately follow application of ANCRACK, weed control normally will not be impaired. However, some stunting of growth or even stand reduction may be noted, especially in lighter soils. Cool soil temperatures may accentuate this condition. With the return of good growing conditions, the stunted plants usually will recover and maintain a normal growth pattern.

Note 4. ANCRACK works best in moist soils. If irrigation is used, best results are obtained when water is applied



If symptoms of poisoning develop, send for a physician and treat as in FIRST AID, above. Discard contaminated clothing and shoes, or clean them thoroughly before re-use. Staining may result if contaminated articles are allowed to contact carpets, upholstery, etc.

NOTE TO ATTENDING PHYSICIAN: Active ingredient is a metabolic stimulant. Treat symptomatically.

CONTAINER DISPOSAL: Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Keep out of smoke.

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. **ALL SUCH RISKS ARE ASSUMED BY THE BUYER.**

Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Ansul makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Ansul or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.

GENERAL DIRECTIONS: Liquid ANCRACK is a selective herbicide combination for the control of the following annual grasses and broad-leaf weeds:

ANNUAL GRASSES: Foxtail, crabgrass, barnyardgrass, goosegrass, stinkgrass, sandbur, watergrass, spangletop, windmillgrass, cupgrass and johnsongrass (from seed).

just as the peanuts begin to crack the ground, or "cracking-time". This is normally from 5 to 7 days after planting. However, application can be made from planting time up to the time the majority of the peanut plants are in the "cracking-stage".

RECOMMENDED DOSAGE RATE FOR PEANUTS:

Use ANCRACK at the rate of 6 quarts in 20 to 40 gallons of water per acre as an overall spray (broadcast application). Treatment may be used over the entire area or in bands as desired. The soil should not be disturbed after planting when the "cracking-time" treatment is used.

Amount of ANCRACK Herbicide Per Acre to Use for Band Treatment of Peanuts

Width of Band	Row Spacing		
	24 inches	28 inches	30 inches
12"	6 pints	5 pints	4 1/4 pints
14"	7 pints	6 pints	5 1/2 pints

Width of Band	Row Spacing		
	32 inches	34 inches	36 inches
12"	4 1/2 pints	4 1/4 pints	4 pints
14"	5 1/4 pints	5 pints	4 1/2 pints

Use in 12-20 gallons of water per acre.

Note 1. Plant on a flat or slightly raised band.

Note 2. Space seed in row about 3 inches (Spanish) or 4 inches (other varieties) and plant at least 2 inches deep.

Note 3. May be used in the same manner following the use of a preplant incorporated herbicide.

Do not allow spray to contact Ornamentals or Crop Plants not recommended on label as injury may result.

CONTAINER STORAGE AND HANDLING:

ANCRACK is normally a solution but it may crystallize in the cold (32-40°F.). Gentle warming (75-125°F.) will redissolve any crystals. Mild agitation, such as rolling, shaking or stirring before use is recommended.

Avoid contamination of foodstuffs, feeds, fertilizers, seeds, plants and pesticides. Store in a cool dry location.

13 gallons of water per 28 inch row.

Note 1. In the black soil areas of western Indiana, southern Minnesota, northern Iowa and Illinois, the dosage of ANCRACK suggested in the chart above should be increased by approximately one-third.

Note 2. Soybean seed should be well covered with soil. Do not use on shallow planted or exposed seed.

Note 3. If heavy rains immediately follow application of ANCRACK, weed control normally will not be impaired. However, some stunting of growth or even stand reduction may be noted, especially in lighter soils. Cool soil temperatures may accentuate this condition. With the return of good growing conditions, the stunted plants usually will recover and maintain a normal growth pattern.

Note 4. ANCRACK works best in moist soils. If irrigation is used, best results are obtained when water is applied after applying the chemical mixture.

Note 5. May be used in the same manner following the use of a preplant incorporated herbicide.

ANCRACK Herbicide is manufactured by The Ansul Company, Marinette, Wisconsin 54143.

X . 81. ANSUL, ANCRACK are registered trademarks of The Ansul Company.

EPA Reg. No. 6308-73

Net Contents 5 Gallons

Form No.

Jan 9, 1973

6308-73



THE ANSUL COMPANY, MARINETTE WISCONSIN

ANSUL LABEL