

CAUTION: Keep Out of the Reach of Children

CAUTION: Humilian diswallow it Applicant at this kin. Avoid preathing apply med. Vrain hands after using Avoid storage near feed or food products. Keep child en and domestic an mark off the ded areas until this material has been

ANTIDOTE If taken internally, induce voniting and call physician at onle

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. WARRANTY - CONDITION OF SALE. DIRECTIONS FOR USE of this product are based on field use and tests betieved reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Setter of this product, such things as crop injury, ineffectiveness or

other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Ansul makes no other warr inties, express or implied, including FITNESS or MERCHAUTABILITY. In no case shall Ansul or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by

The Ansul Company and is accepted as such by the Buyer.

GENERAL INFORMATION: ANSAR 529 H.C. Herbicide (Western Formulation) is useful for selective post emergent weed control, particularly grassy weeds. This ANSAR 529 H.C. Herbicide (Western Formulation) has been specifically formulated for use in the arid and semi-arid regions of the Western United States. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds. The phytotoxic properties are quickly inactivated on contact with the soil.

MIXING INSTRUCTIONS: ANSAR 529 H.C. Herbicide (Western Formulation) is completely water soluble. Fill the spray tank about half full with water and add the required amount of herbicide. Finish filling the tank with water using mild agitation. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period. Although ANSAR 529 H.C. Herbicide (Western Formulation) is only moderately corrosive, do not use in galvanized or aluminum equipment.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible and discard in a safe place

Do not feed treated foliage to livestock or graze treated areas. Do not contaminate waters used for domestic consump tion, or by animals, wildlife and aquatic life, or for irrigation purposes.

DIRECTIONS FOR USE:

COTTON (Arizona - California). ANSAR 529 H.C. Herbicide (Western Formulation) is useful as a directed application in cotton for post-emergent control of weeds and grasses such as johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, sandbur, crabgrass and puncture vine. It should be applied at the rate of 173 galton per acre. Mix 7 pints of ANSAR 529 H.C. Herbicide (Western Formulation) in 100 gallons of water. Do not add any surfactant

to the spray solution. Apply spray solution at 40 gallons per treated acre. Best results are obtained on young actively growing weeds. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre of cotton. Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage. Where practical, withhold irrigation for approximately ten (10) days. If regrowth occurs, make a second application 1 to 3 weeks after the first

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

Apply only when cotton is 3 inches high to first bloom. Do not apply after first bloom. Do not feed treated foliage to livestock or graze treated areas.

Preplant Application: If planting of cotton has been delayed and weeds have emerged, use a preplant application Cotton may be planted immediately following the preplant application.

CITRUS - BEARING AND NON-BEARING (Except Florida): ANSAR 529 H.C. Herbicide (Western Formular on) is useful as a directed application in citrus orchards, such as orange, grapefruit, tangerine, iemon and lime orchards. It should be applied at the rate of 1.3 to 2.3 gailons per acre-

Mix ANSAR 529 H.C. Herbicide (Western Formulation) at the rate of 1/3 gallon in 75 gallons of water. Apply as a directed spray in interspaces and around base of trees. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year. Do not allow spray solution to contact fruit, leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees

As a spot treatment mix ANSAR 529 H.C. Herbicide (Western Formulation) at the rate of 1/3 gallon in 100 gallons of water. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year. Do not allow spray solution to contact fruit, leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees

AGRICULTURAL PLANTINGS: ANSAR 529 H.C. Herbicide (Western Formulation) is useful as a directed application in non-bearing vineyards and in non-bearing faciduous fruit and nut orchards, such as apple, peach, pear plum, prune, appricot, cherry, atmond and walnut orchards. It should be applied at the rate of 1/3 to 2/3 gailons

per acre.
Mix ANSAR 529 H.C. Herbicide (Western Formulation) at the rate of 1/3 gallon in 50 to 100 gallons of water. Apply as a directed spray in interspaces and around base of trees or vines, Spray unwanted vegetation to just short of run-off If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact leaves, stems or bank. Use a shield, if necessary, for nursery plantings or young trees. Do not apply around trees or vines from which fruit will be harvested within one year of treatment. NON-CROP: ANSAR 529 H.C. Herbicide (Western Formulation) is useful for control of johnsongrass, nutsedge, dallisgrass, cocklebur, barrenda or crabgrass, sandbur, ragweed, puncture vine, tules and certain other weeds on

drainage dischbanks, rights-of-way, fence rows, storage yards and similar non-crop areas.

Mix ANSAR 529 H.C. Herbicide (Western Formulation) at a rate of 3 to 6 pints in 100 gallons of water. Spray unwanted vegetation sufficiently to provide complete coverage. Use spray equipment that gives good low volume

coverage. If regrowth occurs, reapply as required.

LAWN AND ORNAMENTAL TURF ANSAR 529 HC Herbicide (Western Formulation) is useful for control of dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, chickweed and wood sorrel with little or no injury to tolerant lawn grasses. On new lawns, do not treat until after three mowings. Tolerant grasses may be temporarily discolored Zoysiagrass, bluegrasses and bermudagriss are quite tolerant. Do not use on St. Augustine or centipede.

Mow lawns 1 to 1 1/2 inches high before treatment. Mix 1 1/3 fluid ounces (8 teaspoons) of ANSAR 529 H.C. Herbicide (Western Formulation) in 2.1.2 gallons of water and apply to an area of 1000 sq. ft. pray thoroughly to wet all undestrable plants. Repeat applications, 10 to 14 days apart, may be needed for good, antro-

X II ANSAR, ANSUL are required trademarks of The Ansul Company

Net Conte' & 5 Gallons Form No.

EPA Reg. No. 6308 77

0]

THE ANSUL COMPANY, MARINETTE, WISCONSIN