

STATE REGISTRATION: FLORIDA

FLORIDA LAW prohibits the use of this material in a broadcast manner.

CAUTION: Keep Out of the Reach of Children

CAUTION: Harmful if swallowed. Avoid contact with skin, Avoid breathing spray mist, Wash hands after using, Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been

ANTIDOTE: If taken internally, induce vomiting and call physician at once. READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

WARRANTY - CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Ansul makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Ansul or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.

GENERAL INFORMATION: ANSAR 529 H.C. Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its phytotoxic properties are quickly inactivated on contact with soil. It is a combination of herbicide and surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young actively growing weeds at air temperatures above 70°F.

MIXING INSTRUCTIONS: ANSAR 529 H.C. Herbicide is completely water soluble, Fill the spray equipment tank about half full with water and add the required amount of herbicide with agitation, Finish filling the tank with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

Do not contaminate waters used for domestic consumption, or by enimals, wildlife and equatic life, or for irrigation purposes. Do not feed treated foliage to livestock or graze treated areas.

COTTON: ANSAR 529 H.C. Herbicide is useful as a directed application in cotton for post-emergent control of weeds and grasses, such as johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, crabgrass, barnyardgrass, sandbur

Mix 3 quarts of ANSAR 529 H.C. Herbicide in 90 gallons of water. Do not add any surfactant to the spray solution. Weed control is most effective at temperatures above 70°F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre.

Band Application: Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the LOTE in Fillage, but to give good coverage of the weeds and grasses. If regrowth occurs, make a second application application application and a reddish discoloration of the cotton leaf may occasionally grow following directed band application, but the cotton plant will develop normally.

Make band applications only when cotton is 3 inches high to first bloom. Do not apply after first bloom.

Preplant Application: If planting of cotton has been delayed and weeds have emerged, use one preplant application. Apply band application just ahead of planter. Cotton may be planted immediately following the preplant application. BEARING CITRUS: ANSAR 529 H.C. Herbicide is useful as a directed application in citrus orchards, such as orange, grapefight, tangerine, lemon and lime orchards. It should be applied at the rate of 1/3 gallon per acre. Mix ANSAR 529 H.C. Herbicide at the rate of 1/3 gallon in 50 gallons of water. Apply as a directed spray in interspaces and around base of trees. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact fruit, leaves, stems or bark. Use a shield, if necessary, for nursery plantings or

AGRICULTURAL PLANTINGS: ANSAR 529 H.C. Herbicide is useful as a directed application in non-bearing vineyards and in non-bearing deciduous fruit, nut and citrus orchards, such as apple, peach, pear, plum, prune, apricot, cherry, almond, walnut, orange, grapefruit, tangerine, lemon and lime orchards.

Mix ANSAR 529 H.C. Herbicide at the rate of 1/3 gallon in 50 gallons of water. Apply as a directed spray around base of trees or vines. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact leaves, stems or bark. Use a shield, if necessary, for nursery plantings of young trees. Do not apply around trees or vines from which fruit will be harvested within one year of treatment, NON-CROP: ANSAR 529 H.C. Herbicide is useful as a spot applied herbicide for control of johnsongrass, barnyardgrass, nutsedge, crabgrass, dallisgrass, cocklebur, ragweed, sandbur, puncture vine and certain other weeds on

drainage ditchbanks, rights-of-way, fence rows, storage yards and similar non-crop areas. Mix ANSAR 529 H.C. Herbicide it a rate of 6 to 12 pints in 100 gallons of water. Make spot applications on unwanted vegetation, spray thoroughly, but just short of run-off. Use spray equipment that gives good low volume coverage. If regrowth occurs, reapply as required,

LAWN AND ORNAMENTAL TURF: ANSAR 529 H.C. Herbicide is useful for control of dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, chickweed and wood sorrel with little or no injury to tolerant lawn grasses. On new lawns, do not treat until after three mowings. Tolerant grasses may be temporarily discolored. Zoysia grasses, blue grasses and bermuda grasses are quite tolerant. Do not use on St. Augustine or centipede grasses.

Mow lawns 1 to 1 1/2 inches high before treatment. Mix 1 1/3 fluid ounces (8 teaspoons) of ANSAR 529 H.C. Herbicide in 2 1/2 gallons of water. This will make enough spray material when used as a spot application to cover a total of 1000 sq. ft. Spot spray thoroughly to wet all undesirable plants. Repeat applications, 10 to 14 days apart, may be needed for good control.

ANSAR 529 H.C. Herbicide is manufactured by The Ansul Company, Marinette, Wisconsin 54143.

X . 31, ANSAR, ANSUL are registered trademarks of The Ansul Company.

Net Contents 5 Gallons

EPA Reg. No. 6308-85

THE ANSUL COMPANY, MARINETTE, WISCONSIN

NEW USE SECTION "Bearing Citrus"


BEARING CITRUS: ANSAR 529 H.C. Herbicide is useful as a directed application in citrus orchards, such as orange, grapefruit, tangerine, lemon and lime orchards. It should be applied at the rate of 1/3 gallon per acre. Mix ANSAR 529 H.C. Herbicide at the rate of 1/3 gallon in 50 gallons of water. Apply as a directed spray in interspaces and around base of trees. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact fruit, leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees.

#3 NEW DISCLAIMER STATEMENT

WARRANTY - CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Ansul makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Ansul or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.



STATE REGISTRATION: FLORIDA FLORIDA LAW prohibits the use of this miterial in a broadcast manner, CAUTION: Keep Out of the Reach of Children CAUTION: Harmful if swallowed. Avoid contact with skin, Avoid breathing spray mist, Wash hands after using, Avoid

washed into the soil. protongrid period. DIRECTIONS FOR USE:

and puncture vine Young tore

Form No.

storage near feed or food products. Keep children and domestic animals off treated areas until this material has been

ANTIDOTE If taken internally, induce vomiting and call physician at once, READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

WARRANTY - CONDITION OF SALE. DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as cropinjury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to an the directions for use as modified by the above. Ansul makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Ansul or the Seiler be liable for consequential special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.

GENERAL INFORMATION: ANSAR '29 H.C. Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its phytotoxic properties are quickly inactivated on contact with soil. It is a combination of herbicide and surfactant. It is unnecessary to add any other surfactant to the spray solution, Best results are obtained on young actively growing weeds at air temperatures above 70°F.

MIXING INSTRUCTIONS: ANSAR 529 H.C. Herbicide is completely water soluble, Fill the spray equipment tank about haif full with water and iidd the required amount of herbicide with agitation. Finish filling the tank with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place

Do not contaminate waters used for domestic consumption, or by animals, wildlife and equatic life, or for irrigation purposes. Do not feed treated foliage to livestock or graze treated areas.

COTTON: ANSAR 529 H.C. Herbicide is useful as a directed application in cotton for post-emergent control of wends and grasses, such as johnsonglass, nutsedge, dallisgrass, conklebur, ragweed, crabgrass, barnyardgrass, sandbur

Mix 3 quarts of ANSAR 529 H.C. Herbicide in 90 galions of water. Do not add any surfactant to the spray solution. Weed control is most effective at temperatures etway 709E. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per inclu-

Band Application: Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage, but to give good coverage of the weeds and grasses. If regrowth occurs, make a second application about 1 to 3 weeks after the first. Slight burning and a reddish discoloration of the cotton leaf may occusionally occur following directed band application, but the cotton plant will develop normally.

Make band applications only when cotton is 3 inches high to first bloom. Do not apply after first bloom.

Preplant Application: If planting of cotton has been delayed and weeds have emerged, use one preplant application. Apply band application just ahead of planter. Cotton may be planted immediately following the preplant application BEARING CITRUS: ANSAR 529 H.C. Herbicide is useful as a directed application in citrus orchards, such as orange, gradefulit, tungerme, lemon and time orchards. It should be applied at the rate of 1/3 gallon per acre. Mix ANSAR 529 H.C. Herbicide at the rate of 1/3 gallon in 50 gallons of water, Apply as a directed spray in interspaces and around base of treps. Spray unwanted vegetation to just short of run-off, If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spruy solution to contict fruit, leaves, stems or bark. Use a shield, if necessary, for nursery plantings or

AGRICULTURAL PLANTINGS: ANSAR 529 H.C. Herbicide is useful as a directed application in non-bearing ver-yards and in non-bearing deciduous fruit, nut and citrus orchards, such as apple, peach, pear, plum, prune, apricut, Carry, almond, walnut, or inge, grapefruit, talgerine, lemon and lime orchards.

Mix ANSAR 529 H.C. Herbicide at the rate of 1/3 gallon in 50 gallons of water. Apply as a directed spray around base of trees or vines. Sprily unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact leaves, stems or bark. Use a shield, if necessary, for nursery plantings of young trees. Do not apply around trees or vines from which fruit will be harvested within one year of treatment. NON-CROP. ANSAR 529 H.C. He bicide is useful as a spot applied herbicide for control of johnsongrass, barnyard grass, nutsedge, crabgrass, dallisgrass, cricklebur, ragweed, sandbur, puncture vine and certain other weeds on

drainage ditchbunks, rights of way, fence rows, storage yards and similar non-crop areas. Mix ANSAR 529 H.C. Herbicide it a rate of 6 to 12 pints in 100 gallons of water. Make spot applications on unwarded ingetation, spray thoroughly, but just short of run-off. Use spray equipment that gives good low we ome (roundless). If regrowth occurs, reapply as required,

LAWN AND ORNAMENTAL TURF: ANSAR 529 H.C. Herbicide is useful for control of dallisgrass, sandbur, bab agrass, putsingly, crubyrass, chicksweed and wood sorrel with little or no injury to tolerant lawn grasses. On new lawns, do nut treat until after three mowings. Tolerant grasses may be temporarily discolored. Zoysia grasses, bluegrasses and bermuda grasses are quite tolerant. Do not use on St. Augustine or centipede grassis,

Mow lawns 1 to 1 1/2 inches high before treatmant, Mix 1 1/3 fluid ounces (8 teaspoons) of ANSAR 529 H.C. Herbicide in 2 1/2 gallons of water. This will make enough spray material when used as a spot application to cover a total of 1000 sq. ft. Spot spray thoroughly to wet all undesirable plants, Repeat applications, 10 to 14 days apart, muy be needed for good control.

ANSAR 529 H.C. Herbicide is manufactured by The Ansul Company, Marinette, Wisconsin 54143.

. . ANSAR, ANSUL ire registered trademarks of The Ansul Company.

Net Contents 5 Gallons

EPA Reg. No. 6308-85

THE ANSUL COMPANY, MARINETTE, WISCONSIN

$= 2 \qquad \qquad \sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{i=1}^{n} \sum_{i=1}^{n} \sum_{i=1}^{n} \sum_{j=1}^$

a contract of the second second a Vising Salar and a state of the second state of the second state of the second state of the second state of the and a second المنافع المنافع المنافع المنافع المنافع والمنافع و a construction of the second second Since and and a second se do na exace a a sister service

To not allow splat second a to concert place as a second pro-Use a shield, is necessary, for marsery practings of young thees.

#3 NEW DISCLAIMER STATEMENT

.

WARRANN - CONTITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. it is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, ineffectlyeness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED ON THE SUVER.

Assul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purjoses referred to in the directions for use as modified by the above. Ansal makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Ansul or the Seller be liable for consequential, special or indirect damages. resulting from the ase or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.