

Ansar 170 H.C.

Herbicide

HIGH CONCENTRATE

MSMA Liquid

For Selective
Post-Emergent
Weed Control

ACTIVE INGREDIENTS:

Monosodium Acid
Methanearsonate* 58.39%


INERT INGREDIENTS: 41.61%

Total Arsenic (as elemental)
all in water soluble
form 27.01%

*Product contains 8.00 lbs.
MSMA per gallon.

CAUTION:
Keep out of the
reach of children

Read entire label
before using
this product



STATE REGISTRATION: FLORIDA

FLORIDA LAW prohibits the use of this material in a broadcast manner.

CAUTION: Keep Out of the Reach of Children

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

WARRANTY - CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Ansul makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Ansul or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.

GENERAL INFORMATION: ANSAR 170 H.C. Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its activity is enhanced by addition of a suitable surfactant to the spray solution. It is most effective on young, actively growing weeds at air temperatures above 70°F. Its phytotoxic properties are quickly inactivated on contact with the soil.

MIXING INSTRUCTIONS: ANSAR 170 H.C. Herbicide is completely water soluble. Fill the spray equipment reservoir about half full with water and add the required amount of herbicide and surfactant with agitation. Finish filling the reservoir with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes. Do not feed treated foliage to livestock or graze treated areas.

DIRECTIONS FOR USE:

COTTON: ANSAR 170 H.C. Herbicide is useful as a directed application in cotton for the post-emergent control of weeds and grasses such as johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, crabgrass, barnyardgrass, sandbur and puncture vine.

Add 5 pints of ANSAR 170 H.C. Herbicide and 1 to 2 quarts of a suitable surfactant to 100 gallons of water and mix thoroughly. Weed control is most effective at temperatures above 70°F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre.

Band Application: Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage, but to give good coverage of the weeds and grasses. If regrowth occurs, make a second application about 1 to 3 weeks after the first. Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following directed band application, but the cotton plant will develop normally.

Make band applications only when cotton is 3 inches high to first bloom. Do not apply after first bloom.

NON-CROP: ANSAR 170 H.C. Herbicide is useful as a spot applied herbicide for control of johnsongrass, barnyardgrass, nutsedge, crabgrass, dallisgrass, cocklebur, ragweed, sandbur, puncture vine and certain other weeds on drainage ditchbanks, rights-of-way, fence rows, storage yards and similar non-crop areas.

Mix ANSAR 170 H.C. Herbicide at a rate of 2 to 5 pints, plus 1 to 2 quarts of a suitable surfactant in 100 gallons of water. Make spot applications on unwanted vegetation, spraying thoroughly, but to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

LAWN AND ORNAMENTAL TURF: ANSAR 170 H.C. Herbicide is useful for control of dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, chickweed and wood sorrel with little or no injury to tolerant lawn grasses. On new lawns, do not treat until after three mowings. Tolerant grasses may be temporarily discolored. Zoysia grasses, bluegrasses, and bermuda grasses are quite tolerant. Do not use on St. Augustine or centipede grasses.

Mow lawns 1 to 1½ inches high before treatment. Mix 1 fluid ounce (2 tablespoons) of ANSAR 170 H.C. Herbicide and ½ fluid ounce (1 tablespoon) of a suitable surfactant in 5 gallons of water. This will make enough spray material when used as a spot application to cover a total of 1000 sq. ft. Spot spray thoroughly to wet all undesirable plants. Repeat applications, 10 to 14 days apart, may be needed for good control.


ANSAR 170 H.C. Herbicide is manufactured by The Ansul Company, Marinette, Wisconsin 54143.

X, CC, ANSAR, ANSUL are registered trademarks of The Ansul Company.

Net Contents 5 Gallons

EPA Reg. No. 6308-86

Form No.



THE ANSUL COMPANY,
MARINETTE, WISCONSIN

SPECIMEN LABEL

Herbicide

ANSAR 170 H.C. HERBICIDE

MSMA Liquid

For Selective Post-Emergent Weed Control

ACTIVE INGREDIENTS:

Monosodium Acid Methanearsonate* 58.39%


INERT INGREDIENTS: 41.61%

Total Arsenic (as elemental) all in water soluble form 27.01%

*Product contains 8.00 lbs. MSMA per gallon.

CAUTION: Keep out of the reach of children

Read entire label before using this product



SPECIMEN LABEL

STATE REGISTRATION: FLORIDA

FLORIDA LAW ENFORCEMENT AGENCY

ANSAR 170 H.C. HERBICIDE

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near food or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

WARRANTY - CONDITION OF SALE: DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however not possible to minimize all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of The Ansul Company or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result. ALL SUCH RISKS ARE ASSUMED BY THE BUYER.

Ansul warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use as modified by the above. Ansul makes no other warranties, express or implied, including FITNESS or MERCHANTABILITY. In no case shall Ansul or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by The Ansul Company and is accepted as such by the Buyer.

GENERAL INFORMATION: ANSAR 170 H.C. Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its activity is enhanced by addition of a suitable surfactant to the spray solution. It is most effective on young, actively growing weeds at air temperatures above 70°F. Its phytotoxic properties are quickly inactivated on contact with the soil.

MIXING INSTRUCTIONS: ANSAR 170 H.C. Herbicide is completely water soluble. Fill the spray equipment reservoir about half full with water and add the required amount of herbicide and surfactant with agitation. Finish filling the reservoir with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes. Do not feed treated foliage to livestock or graze treated areas.

DIRECTIONS FOR USE:

COTTON: ANSAR 170 H.C. Herbicide is useful as a directed application in cotton for the post-emergent control of weeds and grasses such as johnsongrass, nuttallgrass, dallisgrass, cocklebur, ragweed, crabgrass, barnyardgrass, sandbur and puncture vine.

Add 5 pints of ANSAR 170 H.C. Herbicide and 1 to 2 quarts of suitable surfactant to 100 gallons of water and mix thoroughly. Weed control is most effective at temperatures above 70°F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre.

Band Application: Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage, but to give good coverage of the weeds and grasses. If regrowth occurs, make a second application about 1 to 3 weeks after the first. Slight curling and a reddish discoloration of the cotton leaf may occasionally occur following directed band application, but the cotton plant will develop normally.

Make band applications only when cotton is 3 inches high to first bloom. Do not apply after first bloom.

NON-CROP: ANSAR 170 H.C. Herbicide is useful as a spot application for control of johnsongrass, barnyardgrass, nuttallgrass, crabgrass, dallisgrass, cocklebur, ragweed, sandbur, puncture vine and certain other weeds on garage ditchbanks, rights of way, fence rows, storage yard and other non-crop areas.

Mix ANSAR 170 H.C. Herbicide at a rate of 2 to 5 pints per acre with 2 quarts of a suitable surfactant in 100 gallons of water. Make spot applications on unwanted weeds, spraying thoroughly, but to just short of run off. Any spray equipment that gives good coverage may be used for spot applications, applying as required.

LAWN AND ORNAMENTAL TURF: ANSAR 170 H.C. Herbicide is useful for control of dallisgrass, sandbur, bahiagrass, nuttallgrass, crabgrass, chickweed and 1 wood sorrel with little or no injury to tolerant lawn grasses. On new lawns, do not treat until after three mowings. Tolerant plants may be temporarily discolored. Zoysia grasses, bluegrasses and bermuda grasses are quite tolerant. Do not apply to St. Augustine or centipede grasses.

Mow lawns 1 to 1 1/2 inches high before treatment. Mix 1 fluid ounce (2 tablespoons) of ANSAR 170 H.C. Herbicide and 1 fluid ounce (1/2 cup) of a suitable surfactant in 5 gallons of water. This will make enough spray material when used as a spot application to cover a total of 1600 sq. ft. Spot spray thoroughly to wet all undesirable plants. Repeat applications 10 to 14 days apart, may be used for continued control.

ANSAR 170 H.C. Herbicide is manufactured by The Ansul Company, Marinette, Wisconsin 54143.

X, O, ANSAR, ANSUL being registered trademarks of The Ansul Company.

Net Contents 5 Gallons Form No.

EPA Reg. No. 6308-86

THE ANSUL COMPANY, MARINETTE, WISCONSIN