

CAUTION!

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

FSN 6840-753-4972 RODENTICIDE, ANTICOAGULANT, UNIVERSAL, CONCENTRATE

NET WEIGHT 1 LB.

LEVEL A

ACTIVE INGREDIENTS:

Di-(2-ethyl-1,3-indandione (acid equivalent)) 0.3%

NEPT INGREDIENTS:

Tetrasodium ethylenediamine tetracetate 3.0%

Sodium Benzoate 3.0%

Sugar 93.7%

TOTAL 100.0%

Manufactured by
TYLER PRODUCTS
4525 - 5th St. N.E.
Puyallup, Wash. 98371

DIRECTIONS

Mix can be used for Either Liquid or Dry Cereal Baits.
LIQUID BAITS - Dissolve contents of one pound can in 4 gallons of water, or approximately 2 tablespoons in one quart of water.
CEREAL BAITS - Thoroughly mix one part (by weight) of rodenticide concentrate with 10 parts of bait material, such as corn meal, breakfast cereal, oats, or suitable combinations.

PLACEMENT - Place baits in locations where rats and mice feed, water or travel. Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable. Bait should be placed in locations not accessible to children, pets and domestic animals, or in tamper-proof boxes. Contaminated or spoiled baits should be replaced promptly. Baiting should continue until all signs of feeding have stopped. Where a continuous source of infestation is present, permanent bait stations should be established and replenished as required.

FOR RATS - Use at least four ounces of bait (or pint liquid dispenser) per station. An uninterrupted supply of bait should be maintained for at least 10 days.

FOR MICE - Use tablespoon amounts (1/4 to 1/2 ounce of bait or pint liquid dispenser) at 8 to 12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

CAUTION - Keep away from humans, domestic animals, and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K, combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of 5-hydroxyvitamin K.

EPA Reg. No. 5311-7

CUSTOMER *General Can Co*
LAYOUT *094 611 011*
COLORS *BLACK*
REV 11 24
ENG DEF

8132-4-3-68
20X4 COASTER DOUBLET
SEE 307X315
ARTIST *9104*

ACCEPTED