

# AIR-FLOATED DUSTING SULFUR

1. ACTIVE INGREDIENT: SULFUR . . . . . 93%
2. INERT INGREDIENTS: . . . . . 7%
3. FINESS: NOT LESS THAN 93% PASSING 325 MESH

## Directions For Use

4. Yellow Jacket Dusting Sulfur is formulated for use in mixtures or alone as a dust when a 325 mesh free flowing dust is required.  
**IMPORTANT:** For best results, apply when air is calm, just often enough to keep foliage and fruit covered with a light uniform coating.  
**CAUTION:** Do not use sulfur within two weeks of an oil spray.  
Do not use sulfur on apples when temperature is high.  
Do not use sulfur on cucurbit, 'cucumber, cantaloupe, melon, squash' or tung.  
To do so may result in severe injury to foliage and fruit.
5. Peanuts - *Cercospora* Leaf Spot - Use 8.0 to 24.0 pounds as dust per acre. Begin when peanuts are 40 days old or when first spots appear on lower leaves. Repeat at 10 to 14 day intervals and after rains.
6. Sulfur may burn foliage when temperatures are excessively high.  
Do not apply when foliage is damp or wet, as burning may result.

***DANGER: Keep Out of Reach of Children***

Manufactured by  
**GEORGIA GULF SULPHUR CO.**  
BAINBRIDGE, GEORGIA

USA Reg 6325-3

