

MICRO FINE WETTABLE SULFUR

EPA REG. NO. 6325-10

STORAGE AND DISPOSAL (12/17)

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store unused product in original container only in cool, dry area out of reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.

ACCEPTED

DEC 16 1983

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, the pesticide  
registered with EPA Reg. No. 6325-10

## RECOMMENDATIONS

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing and number of applications needed will vary with local conditions.

All dosages given on this label are for dilute sprays (1X concentrations). For use in concentrate sprays, increase the

**IMPORTANT:** Use the higher dosage during periods of frequent and heavy rainfall. Spray often enough to keep susceptible growth covered. This may require weekly or bi-weekly applications during periods of rapid growth and frequent rainfall. When using Lead Arsenate with Micro-Fine Wettable Sulfur always add the same amount of Spray Lime, or other safener, as would be used when applying Lead Arsenate without sulfur. When using Zinc Sulfate and Spray Lime as a safener on peaches, always add Zinc Sulfate to the Spray tank first, followed by Spray Lime, Lead Arsenate and sulfur in order given.

**Apple, Pear: Powdery Mildew, Scab—** Use 6 to 8 lbs. per 100 gals. of water in the pre-bloom, bloom and petal fall sprays. Use 4 to 6 lbs. in the cover sprays.

**Cherry: Brown Rot, Leaf Spot—** Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, petal fall, shuck, cover and post-harvest sprays.

**Citrus: Rust Mite—** Use 5 lbs. per 100 gals. of water. Begin applications at first signs of damage and repeat as necessary. Rust mites should be controlled at all times of the year in order to prevent russeting of fruit and greasy spot on leaves. **Scale Crawlers—** Add 5 to 8 lbs. to each 100 gals. of tank strength fine sulfur solution.

**Peach: Brown Rot, Scab—** Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, shuck, cover and pre-harvest sprays.

**Plum, Prune: Brown Rot, Leaf Spot—** Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, shuck, cover and pre-harvest sprays.

**Beans, Peas: Powdery Mildew, Rust—** Use 6 lbs. per acre in sufficient water for thorough coverage. Repeat applications in 7 to 14 day intervals as necessary.


**Roses: Black Spot, Powdery Mildew—** Use 6 to 8 lbs. per 100 gals. of water (3 to 4 tablespoons per 1 gal. of water). Make first application prior to first signs of disease and maintain preventive schedule throughout the season, repeating applications every 7 to 14 days. Keep new growth protected.

**Perennials, Shrubs: Powdery Mildew, Rust—** Use 6 to 8 lbs. per 100 gallons of water (3 to 4 teaspoons per 1 gallon of water). Make first application prior to first signs of disease and maintain preventive schedule throughout the season, repeating applications every 7 to 14 days. Keep new growth protected.

## WARRANTY STATEMENT

GEORGIA GULF SULFUR CORPORATION warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the crops specified. There are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

Amount of product by the concentrate factor. For example, if normal use is 3 lbs. per 100 gallons applied to one acre, the 5X concentrate rate will be 25 lbs. per 100 gallons. Spray often enough to keep new growth protected.



BEST DOCUMENT AVAILABLE

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed. Avoid breathing vapors or spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate food or feed products. Keep out of reach of children.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and wildlife. Do not apply directly to lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

**GENERAL CLASSIFICATION**

It is a violation of Federal law to use this label in a manner inconsistent with its labeling.

**RE-ENTRY STATEMENT**

Do not enter treated area without protective clothing and respiratory protection for 24 hours following application.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act. Dispose of empty bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

See & Mac  
replacemnt

**GENERAL INFORMATION**

Apply this product only as specified on this label. Do not use this product for any uses other than those specified on this label.

Micro-Fine **Wettable Sulfur** is formulated for use in water as a spray on fruit, vegetable and ornamental crops for the control of certain fungus diseases and mites. It may be combined in the spray tank with most of the commonly used fungicides and insecticides.

Pour the specific amount of this product on surface of water in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants.

**CAUTION:** Do not use sulfur in combination with oil or within two weeks of an oil spray. Do not use sulfur on apples or other sensitive crops when shade temperatures exceed, or are likely to exceed, 90°F. Do not apply or allow sulfur to drift to Cucurbits (cucumber, cantaloupe, melon, squash) or tung. To do so may result in severe fruit and leaf injury.

MICRO-FINE

**Wettable Sulfur**

For Spraying Fruits  
Vegetables & Ornamentals

ACTIVE INGREDIENT:	
Sulfur .....	90.0%
INERT INGREDIENTS: .....	10.0%
TOTAL .....	100.0%

FINENESS: Not less than 96% passing 325 mesh.

**KEEP OUT OR REACH OF CHILDREN**

**CAUTION**

If on skin, wash immediately with soap and water  
If in eyes, immediately flush eyes with plenty of clear water for at least 15 minutes  
If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

See side Panel For Additional Precautionary Statements.


EPA REG. NO. 6325-10

EPA EST. NO. 6325-GA-1

**GEORGIA GULF SULFUR CORPORATION**

P.O. Box 1165 • Valdosta, GA • 31603-1165

NET CONTENTS 50 LBS.



**BEST DOCUMENT AVAILABLE**