

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid breathing vapors or spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate food or feed products. Keep out of reach of children.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this label in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated area without protective clothing and respiratory protection for 24 hours following application.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Yellow Jacket Wettable Sulfur II is formulated for use in water as a spray on fruit, vegetable and ornamental crops for the control of certain fungus diseases and mites. It may be combined in the spray tank with most of the common fertilizers and pesticides.

Pour the specific amount of this product on surface of water in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants.

CAUTION: Do not use sulfur in combination with oil or within 21-60 days of an oil spray. Do not use sulfur on apples or other sensitive crops when shade temperatures exceed, or are likely to exceed, 90° F. Do not apply or allow sulfur to drift to Cucurbits (cucumber, cantaloupe, melon, squash) or To do so may result in severe fruit and leaf injury. When growing for processing, consult the processor before applying sulfur.

**Yellow Jacket
Wettable Sulfur II**

**For Dusting or Spraying Fruits,
Vegetables & Ornamentals.**

ACTIVE INGREDIENT:

Sulfur 90.0%

INERT INGREDIENTS:

..... 10.0%

TOTAL 100.0%

FINENESS: Not less than 93% passing 325 mesh

KEEP OUT OF REACH OF CHILDREN


CAUTION

If on skin, wash immediately with soap and water.

If in eyes, immediately flush eyes with plenty of clear water for at least 15 minutes.

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

See Side Panel For Additional Precautionary Statements.



VALDOSTA, GEORGIA 31601

EPA EST NO. 6325-GA 1

NET CONTENTS 50 lbs

SEP 6 1983

Under the Federal Insecticide, Fungicide and Rodenticide Act, registered for the pesticide registration under FFA Reg. No. 6325-13

Consult State Extension Service for applications near

This product

All dosages of use in concentration Sulfur II to a use is 5 lbs. per acre will be 25 a dust, use foliage and for new growth

IMPORT

rain

require

frequent

Sulfur

could

Sulfur

to the

in order

Apple, Pear,

water in the

cover sprays

Cherry: Brown

pink, bloom

Citrus: Rust

first signs of

treated at all

greasy spot of

tank strength

Peach: Brown

bloom, shuck

Plum, Prune:

in pink, bloom

Beans, Peas:

water for th

necessary.

Roses: Black

water (3 to 4

first signs of

season, repeat

treated.

Perennials:

gallons of wa

tion prior to

throughout t

growth prote

GEORGIA G

package con

reasonably fit

to directions

no other wa

duct other th

handling or u

mal condition

buyer assum

Yellow Jacket Wettable Sulfur II

For Dusting or Spraying Fruits,
Vegetables & Ornamentals.

ACTIVE INGREDIENT:	
Sulfur	90.0%
INERT INGREDIENTS:	10.0%
TOTAL	100.0%

FINENESS: Not less than 93% passing 325 mesh

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

When using, wear immediately with soap and water.

If in eyes, immediately flush eyes with plenty of clear water for at least 15 minutes.

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

See Side Panel For Additional Precautionary Statements.

GEORGIA GULF SULFUR
EPA REG NO. 6325-13

SEP 6 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the purpose of registration under FIFRA Act, 1972.

VALDOSTA, GEORGIA 31601
EPA EST. NO. 6325-GA-1

NET CONTENTS 50 lbs

RECOMMENDATIONS

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing and number of applications needed will vary with local conditions.

This product may be used as a dust or a spray.

All dosages given on this label are for dilute sprays (1 X concentrations). For use in concentrate sprays, increase the dosage of Yellow Jacket Wettable Sulfur II to apply the equivalent amount per acre. For Example, if normal use is 5 lbs. per 100 gallons of water applied to one acre, the 5X concentrate rate will be 25 lbs. per 100 gallons of water applied to 5 acres. When used as a dust, use the dosage rate omitting the water. Apply to coat the foliage and fruit with a light uniform residue. Spray often enough to keep new growth covered.

IMPORTANT: Use this product during periods of frequent and heavy rain or after a heavy rain to keep susceptible growth covered. This may require several applications during periods of rapid growth and frequent rain. When using Lead Arsenate with Yellow Jacket Wettable Sulfur, always add the same amount of Spray Lime, or other safener, as would be added when applying Lead Arsenate without sulfur. When using Sulfur and Spray Lime as a safener on peaches, always add Zinc to the spray tank first, followed by Spray Lime, Lead Arsenate and Sulfur in order given.

Apple, Pear: Powdery Mildew, Scab — Use 6 to 8 lbs. per 100 gals. of water in the pre-bloom, bloom and petal fall sprays. Use 4 to 6 lbs. in the cover sprays.

Cherry: Brown Rot, Leaf Spot — Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, petal fall, shuck, cover and post-harvest sprays.

Citrus: Rust Mite — Use 5 lbs. per 100 gals. of water. Begin applications at first signs of damage and repeat as necessary. Rust mites should be controlled at all times of the year in order to prevent russetting of fruit and greasy spot on leaves. **Scale Crawlers** — Add 5 to 8 lbs. to each 100 gals. of tank strength lime sulfur solution.

Peach: Brown Rot, Scab — Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, shuck, cover and pre-harvest sprays.

Plum, Prune: Brown Rot, Leaf Spot — Use 6 to 8 lbs. per 100 gals. of water in pink, bloom, shuck, cover and pre harvest sprays.

Beans, Peas: Powdery Mildew, Rust — Use 16 lbs. per acre in sufficient water for thorough coverage. Repeat applications in 7 to 14 day intervals as necessary.

Roses: Black Spot, Powdery Mildew — Use 6 to 8 lbs. per 100 gals. of water (3 to 4 tablespoons per 1 gal. of water). Make first application prior to first signs of disease and maintain preventive schedule throughout the season, repeating applications every 7 to 14 days. Keep new growth protected.

Perennials, Shrubs: Powdery Mildew, Rust — Use 6 to 8 lbs. per 100 gallons of water (3 to 4 teaspoons per 1 gallon of water). Make first application prior to first signs of disease and maintain preventive schedule throughout the season, repeating applications every 7 to 14 days. Keep new growth protected.

WARRANTY STATEMENT

GEORGIA GULF SULFUR warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the crops specified. There are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

Colson Printing Co. 6072