

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

## CAUTION

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed or foodstuffs.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff conditions are likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### RE-ENTRY STATEMENT

Do not enter treated area without protective clothing and respiratory protection for 24 hours following application.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**CAUTION:** Area treated with YELLOW JACKET FLOWABLE SULFUR on \_\_\_\_\_ Do not enter without appropriate protective clothing for 24 hours following application. In case of accidental exposure, see the Statement for Practical Treatment on this label.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store unused product in original container only in cool, dry area out of reach of children and animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

### PRODUCT INFORMATION

YELLOW JACKET FLOWABLE SULFUR is a flowable sulfur formulation that may be applied as a ground or aerial application. Consult your State Agricultural Experiment Station or Extension Specialist for advice in selecting treatments from this label to best fit local conditions.

**USE ONLY ON CROPS LISTED AND ONLY AS SPECIFIED ON THIS LABEL. EXCEPT ON COTTON,** do not apply when temperatures exceed or are likely to exceed 90°F. Do not use sulfur with oil or within 14 days of an oil spray (for citrus 21 days). Do not combine with emulsifiable liquids.

**NOTE:** Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply or allow to drift to apricots, d'Anjou and Comice pears, cranberries, cucurbits (cucumber, cantelopes, melons, squash,) filberts, spinach, tung trees, walnuts or other sensitive plants. Sulfur may burn foliage and fruit during periods of high temperatures and under certain climate conditions.

YELLOW JACKET FLOWABLE SULFUR can be mixed and applied with liquid fertilizer or water.

**DO NOT STOP AGITATION WITH SULFUR IN SPRAY EQUIPMENT. SULFUR IS HIGHLY CORROSIVE, AND EQUIPMENT SHOULD BE CLEANED THOROUGHLY AFTER EACH DAY'S SPRAYING.**



Flowable Application

For Aerial or Ground Application

### ACTIVE INGREDIENT:

Sulfur as S ..... 53.0%

INERT INGREDIENTS: ..... 47.0%

TOTAL ..... 100.0%

This product contains 6 lbs. of Sulfur per gallon.

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

### STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If on skin, wash thoroughly with soap and water.

If in eyes, wash eyes with plenty of clear water for at least 15 minutes.

See side panel for additional precautionary statements.

E.P.A. Reg. No. E325-15  
E.P.A. Est. No. 6325-GA-1

**GEORGIA GULF SULFUR CORPORATION**

P.O. BOX 1165 • VALDOSTA, GA • 31703-1165

NET CONTENTS 5 Gallons  
**ACCEPTED**

OCT 22 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registration number 6325-15

## RECOMMENDATIONS

### FRUIT TREE CROPS:

Unless otherwise specified for specific crops, dosage rates are given as pints of YELLOW JACKET FLOWABLE SULFUR per 100 gallons water for use in a thorough coverage spray. The total amount of YELLOW JACKET FLOWABLE SULFUR per acre should be 2 to 4 gallons depending on the size of the trees. Because of variations in the types of spray equipment used, a range of low and high rates is listed. For high volume sprayers (output 800 to 1000 gallons spray per acre) use the low rate; for low volume sprayers (output 20 to 300 gallons spray per acre) use the high rate.

### APPLE, PEAR:

Scab: Pre-bloom through calyx sprays ..... 1 1/2 to 3 1/2 pints  
Cover Sprays ..... 1/2 to 2 pints

### Powdery Mildew:

Pre-bloom through calyx sprays ..... 3 pints  
Cover Sprays ..... 2 pints

### CHERRY, PLUM, PRUNE:

Brown Rot, Leafspot: Pink and Bloom Sprays ..... 1 1/2 to 3 1/2 pints  
Petal fall, shuck & cover sprays ..... 1/2 to 1 1/2 pints

### CITRUS:

Rust Mite, Clover Mite: Apply sprays November thru May ..... 1/2 to 4 1/2 pints

**THRIPS** (Early spring Treatment for the reduction of Thrips population): Apply 1 to 2 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.

### PEACH, NECTARINE:

Powdery Mildew, Brown Rot, Scab: Pink and Bloom Sprays ..... 1 1/2 to 3 1/2 pints  
Petal Fall, shuck and cover sprays ..... 1/2 to 1 1/2 pints

Application to mature nectarines may cause discoloration.

### FIELD AND VINE CROPS:

Unless otherwise specified for specific crops, dosage rates are given as pints of YELLOW JACKET FLOWABLE SULFUR per acre. This product should be applied by ground equipment, or aircraft in sufficient water to provide thorough coverage.

**BEANS** (dry, colored, green, snap, lima): Powdery mildew, Rust: Use 4 to 8 pints of YELLOW JACKET FLOWABLE SULFUR per acre in sufficient water for thorough coverage. Begin application at the first sign of disease. Repeat at 7 to 14 day intervals to keep new growth protected.

**COTTON:** Atlantic Mite. 2 pints for early season control and up to 1 gallon for mid to late season control.

**GRAPES:** Powdery Mildew. Use 1 to 2 pints per 100 gallons water, maximum of 8 pints YELLOW JACKET FLOWABLE SULFUR per acre in spring and summer. Make first application when shoots are 6 to 8 inches long, second when shoots are 12 to 15 inches long, and third at about bloom stage. The timing, rate and number of applications will vary with local conditions. Consult State Agricultural Extension Service or State Agricultural Experiment Stations in your region for specific information. Follow the USE PRECAUTIONS given on this label.

**PEANUTS:** For rust use 2 to 4 pints per acre of YELLOW JACKET FLOWABLE SULFUR in enough water for good coverage. Begin 40 to 60 days after planting. Use 7 to 14 day intervals as long as needed to control rust.

**POTATOES:** Powdery Mildew. Use 3 to 4 pints in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3 week intervals.

**SUGAR BEETS:** For control of powdery mildew apply the recommended rate in 10 gallons of water per acre by air at first sign of infection. Use 4 pints per acre in the Pacific Northwest and 8 pints per acre in Arizona. Repeat applications should be made at 10 to 30 day intervals or as necessary throughout the season. Under most disease conditions only one or two applications will be needed.

**Sprinkler Irrigation Water** — Use 1-2 gallons per acre by injecting into sprinkler system at or near the irrigation pump. For center pivot systems apply during one full revolution of the system. For portable or solid set systems, apply during the final 15 minutes of the irrigation period. Do not operate irrigation system after application until deposit has thoroughly dried. Apply the product through irrigation systems containing: 1) anti-siphon and check valves which will prevent water source contamination and overflow of the slurry tank, 2) interlocking controls between the metering device and the water pump to insure simultaneous shut-off.

**STRAWBERRIES:** Powdery Mildew. Use 2 pints. Apply at first sign of infection and repeat at 3 week intervals. For Strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canners' recommendations.

**TOMATOES:** Tomato Russet Mite. Use 1/2 to 1 gallon per acre depending upon size of plants. Use full coverage sprays.

### WARRANTY STATEMENT

GEORGIA GULF SULFUR CORPORATION warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions.

There are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

CPC 019602  
GGC 062565

10F1  
Pm 21  
REG. # 6325-15