

3



**FERRET  
RODENTICIDE**

KILLS NORWAY RATS, ROOF RATS AND HOUSE MICE

**ACTIVE INGREDIENTS**

<b>WARFARIN</b>		
3-alpha-(Acetylbenzyl)-4-hydroxycoumarin	0.025%	
Inert Ingredients	99.975%	
	Total	100.000%

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

**DISPOSAL**  
**DO NOT REUSE EMPTY CONTAINER.**  
**WRAP CONTAINER & PUT**  
**IN TRASH COLLECTION**

Manufactured By  
**FERRET LABORATORIES, INC.**  
Slinger, Wisconsin 53086

EPA Reg. No. 6383-1  
EPA Est. No. 34156-WI-1  
Net Weight One Pound

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS:** For control of Norway Rats, Roof Rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, behind boards, by gnawed openings or in beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

**APPLICATION DIRECTIONS**

**RATS:** slit each inner cellophane bag with sharp instrument and place 1 to 4 bags of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Slit each inner cellophane bag with sharp instrument and apply 1/4 to 1/2 ounce (1/2 level tablespoonful amounts) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**RATS & MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**PRECAUTIONARY STATEMENTS**

**HAZARD TO HUMANS & DOMESTIC ANIMALS**

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K-1 combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdoses of 3-hydroxycoumarin.

**ENVIRONMENTAL HAZARD:**

Keep Out Of Lakes, Streams or Ponds.

**FERRET**

**FERRET**

**ACCEPTED**

JUN 5 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act or other law for the pesticide registered under EPA Reg No. 6383-1