



**SANITATION IN  
FOOD SERVICE  
ESTABLISHMENTS**

Mix 1 oz. SANISOL in 4 gal. hot or cold water for 100 p.p.m. sanitizing solution. Immerse cleaned and rinsed glasses, dishes or utensils for 1 minute. Allow to air dry.  
**RINSE WITH POTABLE WATER BEFORE REUSE.**

**SANITATION IN  
MILK PRODUCTION  
PLANTS**

Mix 1 oz. SANISOL in 4 gal. hot or cold water for 200 p.p.m. sanitizing solution. Immerse cleaned and rinsed equipment and utensils for 2 minutes. Allow to air dry.  
**RINSE WITH POTABLE WATER BEFORE REUSE.**

**GENERAL  
DISINFECTION IN  
FOOD SERVICE  
ESTABLISHMENTS**

Mix 1 oz. SANISOL in 2 gal. hot or cold water for 400 p.p.m. disinfectant solution. Clean and rinse floors, walls, sinks and toilet seats and wipe with solution.

90-626

# **ESCO SANISOL**

## **DEODORANT - GERMICIDE - DISINFECTANT** **A STRONG, ODORLESS SANITIZER**

EPA REG. NO. 6436-1-AA

**ACTIVE INGREDIENTS:** Alkyl\* Dimethyl Benzyl Ammonium Chloride... 10.5%  
\*C<sub>12</sub>-50%, C<sub>14</sub>-40%, C<sub>16</sub>-10%

**ETHANOL** ..... 2.1%

**INERT INGREDIENTS** ..... 87.4%

**DANGER:** Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

**SEE FIRST AID STATEMENTS ON SIDE PANEL**

Rinse empty container thoroughly with water and discard it.

**DILUTE WITH WATER AS DIRECTED**

**DO NOT USE IN SOAP SOLUTIONS**

**NET CONTENTS 1 GALLON**

**Eng-Skell Company**

**SAN FRANCISCO, CA 94103**

**FIRST AID:**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**ACCEPTED**

**ACCEPTED WITH COMMENTS**