Our printing shop will cut this label to the prescribed typing sizes necessary.

FACSIMILS LABEL

LONE STAR 28% EQUIVALENT SUPER HI-PRO-FLY MINERAL - MEDICATED

FOR RUMINANTS ONLY

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle, when used as directed. Aids in the control of stomachworms (Haemonchus, Ostertagis, and Trichostrongylus spp.), nodular worms (Oesophagostomum spp.), largemouth bowelworms (Chabertia spp.) and hookworms (Bunostomum spp.) in cattle when fed in accordance with directions for use. Aids in the prevention of foot rot caused by Sperophorus necrophorus, and in soft tissue lumpy jaw caused by Actinobacillus lignierisi in cattle when fed as directed.

Pesticidally Active Drug Ingredient Phenothiazine Pesticidally Inert Ingredients	•	•	 •	•	•	•	•	•	
Active Drug Ingredient: Ethylene diamine dihydriodide.	•	•	 •	•	•	•	•	•	. 0.01%

CCFPTER	GUARANTEED ANALYSIS Crude Protein not less than
1 -	Crude Fat not less than
SO	Crude Fiber not more than
71 Co	Caldium (Ca) not less than 4.00%
どい	> ZEE Caldium (Ca) not more than
<i>©</i> ∪	三世 Phosphorus (P) not less than 2.50%
	ES MANAGETUM (MR) HOL TERR CHAIL
1	Sage (NaCl) not less than
	NaCl) not more than
•	Vitamin A
	Vitamin D

INGREDIENTS

Plant protein products, processed grain by-products, dried forage products, urea, salt(NaCl), dicalcium phosphate, magnesium oxide, ferrous sulfate, manganese sulfate, zinc sulfate, copper sulfate, cobalt carbonate, vitamin A palmitate (stability improved), irradiated dried yeast (source of vitamin D₂).

Manufactured By
TEXAS FARM PRODUCTS COMPANY
Nacogdoches, Texas 75961

TEXAS PERMIT NO. 38820 EPA Reg. No. 6482-2

50 Pounds (Net Weight)

WARNING: Keep Out Of the Reach Of Children.
Not to be fed to horses, ponies or hogs.
See additional warnings on back panel of this tag.

(20) product)

To be fed at the rate of 1/4 gram of phenothiazine per 100 pounds of body weight per day and not to exceed 2.15 grams per head daily.

Cattle should be given access to 28% Equivalent Super Hi-Pro-Fly several weeks prior to the appearance of adult horn flies in the spring and fed continuously until the flies disapperar inthe fall. 28% Equivalent Super Hi-Pro-Fly will prevent the breeding of horn flies and face flies but have no effect on existing adult fly populations. Since horn flies over-winter in the pupal state, all cattle should be sprayed when the flies first appear. This kills the flies emerging from last year's pupae, and will protect the cattle for about two weeks until the 28% Equivalent Super Hi-Pro-Fly can take effect. Since face flies migrate from farm to farm, it is essential that the larvicide program be carried out on an area basis to assist in the reduction of the adult face fly population.

For best results, animals should be dewormed with a drug such as tramisol or thiabendazole (follow label directions for product use) twice a year, in the spring and fall, for removal of the specified parasites. Continuous low-level treatment, in conjunction with good herd management practices, will control the specified parasites and will help control reinfestation.

WARNING:

Do not administer to lactating dairy animals.

Occasional sensitivity to phenothiazine has been reported in cattle.

Consult a veterinarian before using in severely debilitated animals.

Certain parasites may become resistant to phenothiazine therapy.

