EPA Form 8570-6 (Rev. 5-76)

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

Helent 801-24-92

SIGNATURE OF APPROVING OFFICIAL

- c. Delete "For Ruminants Only." You may substitute "For Cattle Only."
- 3. Provide the following Product Chemistry information:
 - a. In reference to the Confidential Statement of Formula (CSF), the upper and lower certified limits should be based on the pure active ingredients rather than the technical or concentrate.
 - b. Analytical Method: Provide the methods used to analyze for the active ingredients or a full reference for a published method or MRID Number.
 - c. Density: e.g., lbs/cu. ft. for solids.
 - d. Corrosion Characteristics: This information can be noted during the storage stability study.
- 4. Submit five copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca Product Manager 13 Insecticide-Rodenticide Branch Registration Division (H7505C)

Enclosures

JAN 27 1992

 $C_{i,j}(x_{i+1}, x_{i+1}) = 0$

LONE STAR 28% SUPER HI-PRO-MIN WITH RABON ORAL LARVICIDE FOR RUMINANTS ONLY

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT BY WEIGHT Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate INERT INGREDIENTS 99.845% **

TOTAL

100.000%

* Each pound contains 0.70 grams of RABON. ## Refers only to ingredients which are not larvicidal.

GUARANTEED FEED ANLAYSIS Crude Protein, not less than 28.00% (This includes not more than 9.24% equivalent protein from non-protein nitrogen.) Crude Fat, not less than Crude Fiber, not more than Calcium (Ca), not less than Calcium (Ca), not more than Phosphorus (P), not less than Mangnesium (Mg), not less than Salt (NaCl), not less than 14.00% Vitamin A, not less than 9,000 IU/LB Vitamin D3, not less than 1,800 IU/LB

FEED INGREDIENTS

Plant protein products, grain products, processed grain by-products, animal protein products, urea, salt, cane molasses, dicalcium phosphate, magnesium oxide, calcium carbonate, vitamin A acetate, D-activated animal sterol (source of vitamin D3), sodium selenite, manganese sulfate, zinc sulfate, iron sulfate, copper sulfate, ethylenediamine dihydriodide, cobalt carbonate.

> KEEP OUT OF REACH OF CHILDREN CAUTION

See Reverse Of Tay For Additional Precautionary Statements

Net Weight 50 pounds

EPA Reg. No. 6482-__

EPA Est. No. 6492-Tx-1

Manufactured By: TEXAS FARM PRODUCTS COMPANY Nacogdoches, Tx 75963-0009

BEST AVAILABLE CODY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks for protection when handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Feed free choice in covered feeders to beef cattle or dairy cattle. Provide one feeding station per 15 to 20 head of cattle. Animals should consume an average of 1.6 ounces per 100 pounds of body weight per day (1 pound per 1000 pounds). This will supply the recommended average intake of 70 mg. of RABON Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before feeding Lone Star 28% Super HI-PRO-MIN with RABON Oral Larvicide but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas.

If consumption is above 1 pound per 1000 pounds of body weight per day, reduce the number of feeding locations or relocate. If consumption is below 1 pound per 1000 pounds of body weight per day, increase the number of feeding locations. Begin feeding in the spring before flies begin to appear, and continue feeding throughout the summer and the fall.

Lone Star 28% S' or HI-PRO-MIN with RABON Oral Larvicide may be fed up to slaughter and to jactating dairy cows.

Lone Star 28% Super Hi-Pro-Min with RABON Oral Larvicide prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and andund, mattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle marture but in other decaying vegetable matterior silage on the premises. In order to achieve optimum fly control, use this product in conjunction with other good management and sanitation practices:

WARNING: To be fed to ruminants only. Do not feed to horses, mules, colts, ponies or hogs.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. STORAGE: Store in a dry place. CONTAINER DISPOSAL: Completely empty bag into feeding equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

BEST AVAILABLE COPY